

Sky and TELESCOPE

Swain



Indiana University

OCT 24 1957

Library

University of Illinois Observatory

In This Issue:

Swain

N

★
Vol. XVII, No. 1
NOVEMBER, 1957
50 cents
★

American Astronomers Report
A Seeing Compensator
Employing Television
Techniques
A Probable New Meteorite
Fall in Brazil

The Great Aurora
of September 22-23
The Most Massive
Stars Known
Northern and Southern
Star Charts

Celestar[☆]

by fecker



Your key to a new world of enjoyment

An incomparable portable telescope by America's leading manufacturer of astronomical instruments. Although it weighs only 45 pounds its sturdy construction, fork-type mounting and integral electric drive mark it as an instrument of professional quality.

Here's a full-fledged astronomical instrument built to the exacting specifications of a manufacturer of fine telescopes for nearly a century.

Complete with Electric Drive, Slow Motions, Right Ascension and Declination Circles, Heavy Duty Tripod and Finder Telescope.

Magnifications of 35X, 70X, 105X and 210X.

The optics are obviously the best that can be had.

ONLY **\$198.50**
f.o.b. Pittsburgh, Pa.

INDIANA UNIVERSITY LIBRARY

Includes "HANDBOOK of the HEAVENS"

WHEN ORDERING
WRITE DIRECT TO: →

j. w. fecker, inc.

A Subsidiary of AMERICAN OPTICAL COMPANY

6592 HAMILTON AVENUE • PITTSBURGH 6, PA.

EDITORIAL ADVISORY BOARD: *Clement S. Brainin*, Amateur Astronomers Association, New York; *Edward A. Halbach*, Milwaukee Astronomical Society; *Donald H. Menzel*, Harvard College Observatory; *Paul W. Merrill*, Mount Wilson and Palomar Observatories (retired); *Charles H. Smiley*, Ladd Observatory; *Percy W. Witherell*, Bond Astronomical Club.

Vol. XVII, No. 1

NOVEMBER, 1957

CONTENTS

COVER: The University of Illinois Observatory, located on the campus at Urbana, Illinois. It houses a 12-inch Warner and Swasey refractor. The building has recently been enlarged, as described in the adjoining column on this page.

ASTRONOMY AT ILLINOIS	3
AMERICAN ASTRONOMERS REPORT	4
A SEEING COMPENSATOR EMPLOYING TELEVISION TECHNIQUES — John H. DeWitt, Robert H. Hardie, and Carl K. Seyfert	8
A PROBABLE METEORITE FALL IN BRAZIL — Vincent Menezes	10
ARTIFICIAL SATELLITE NO. 1	11
THE GREAT AURORA OF SEPTEMBER 22-23 — James E. McDonald	14
THE MOST MASSIVE STARS KNOWN — Otto Struve	18
DOUBLE STAR STUDIES IN INDONESIA	20
AMATEUR ASTRONOMERS	16
Kansas City Convention Marks League's First Decade	
BOOKS AND THE SKY	31
Light Scattering by Small Particles	
Discoveries and Opinions of Galileo	
The Life of Arthur Stanley Eddington	
Optics: The Science of Vision	
CELESTIAL CALENDAR	46
GLEANINGS FOR ATM's	38
A New Test for Cassegrainian Secondaries	
A Versatile Observing Setup with an 8-inch Reflector	
LETTERS	15
NEWS NOTES	12
OBSERVER'S PAGE	21
The Lunar Eclipse of November 7th	
Deep-Sky Wonders	
September Aurorae	
QUESTIONS	7
SOUTHERN STARS	48
STARS FOR NOVEMBER	49

SKY AND TELESCOPE is published monthly by Sky Publishing Corporation, Harvard College Observatory, 60 Garden St., Cambridge 38, Mass. Second-class mail privileges authorized at Boston, Mass.

Subscription: \$5.00 per year in the United States and possessions; \$9.00 for two years; \$13.00 for three years. Add \$1.00 per year for Canada, Mexico, and all countries of the Pan American Postal Union, making the total subscription \$6.00 for one year; \$11.00 for two years; \$16.00 for three years. Add \$2.00 per year for all other foreign countries, making the total subscription \$7.00 for one year; \$13.00 for two years; \$19.00 for three years. Canadian and foreign remittances should be made in United States currency. Single copy 50 cents. Back numbers, as available, 50 cents each.

All notices of change of address must be sent one month in advance and accompanied by old and new addresses. When sending your renewal order, or writing in regard to your subscription, your current mailing address must be given. Please return our bill form with your renewal payment. Circulation staff: Nancy R. Bolton, manager; Helen B. Sniegiecki; Stacia Strong.

Editorial and advertising offices: Harvard College Observatory, Cambridge 38, Mass. Advertising manager, Henry M. Corrado. Advertising rates are listed in STANDARD RATE AND DATA SERVICE, or sent upon request.

Unsolicited articles and pictures are welcome, bearing adequate return postage, but we cannot guarantee prompt editorial attention, nor are we responsible for the return of unsolicited material. Editorial assistants: Virginia K. Cox; Elizabeth C. Merrylees; William E. Shawcross. The principal articles are indexed in THE READERS' GUIDE TO PERIODICAL LITERATURE.

Astronomy at Illinois

GROUND was broken in 1896 for the University of Illinois Observatory at Urbana, pictured on the front cover of this issue. In the dome is a 12-inch equatorial refractor, made by Warner and Swasey, and there is also a 3-inch combined transit and zenith telescope for student instruction. The observatory has an outpost half a mile to the south, where a 4-inch Ross camera is used in Milky Way research.

This August, just before the American Astronomical Society met at Urbana, an addition was completed that filled in the southwest angle of the original building. It contains a seminar room, offices for the director and a secretary, and a radio astronomy laboratory in the basement. This new wing is at the left in the cover picture, its brickwork yet to mellow to that same rich tone as the original parts of the observatory. The old brick piers in the foreground serve as supports for portable telescopes and other instruments used by astronomy and navigation classes.

Despite its relatively small size and limited observational equipment, the University of Illinois Observatory has exerted a far-reaching influence on modern astronomy.

The first director at Illinois was G. W. Myers, from 1897 to 1900. In the first of these years he analyzed in great detail the light curve of the eclipsing binary Beta Lyrae, deducing the sizes, shapes, densities, and other properties of the two component stars. This was a remarkable achievement 60 years ago, and was praised by Dr. Otto Struve at the Urbana meeting this August, when he gave the Russell lecture on the problems of this famous double star.

In 1903 Joel Stebbins became director, and began a revolution in astronomical photometry. First pushing the light-sensitive selenium cell to its limits for measuring star brightnesses, he later developed the photoelectric cell to obtain extremely precise magnitudes and colors of stars. To this day he has remained in the forefront of this major branch of observational astronomy.

The third director was Robert H. Baker, well known both for his studies of the Milky Way and for his popular writings—his widely used textbook, *Astronomy*, is now in its sixth edition. Dr. Baker served from 1922 to 1951, being succeeded by the present director, George C. McVittie, a British-born astronomer who has made important contributions to the fields of cosmology and relativity.

The last three directors, whose terms span more than half a century, all attended the Urbana meeting, and may be seen together in the photograph by Walter J. Miller, S. J., on page 6.

AMERICAN ASTRONOMERS REPORT

Here are highlights of some papers presented at the 98th meeting of the American Astronomical Society at Urbana, Illinois, in August. Complete abstracts will appear in the *Astronomical Journal*.

Revision of Rowland's Tables

In 1895, the Johns Hopkins physicist, Henry A. Rowland, began publication of an extensive catalogue of the Fraunhofer absorption lines in the spectrum of the sun. These classic tables were revised in 1928 by W. S. Adams and his coworkers at Mount Wilson Observatory.

Now the second revision of Rowland's tables is being prepared jointly by M. Minnaert at the Utrecht Observatory and Charlotte Moore Sitterly at the National Bureau of Standards. The identification of the many thousands of spectral lines, their wave lengths, the multiplet attributions, and the lines' equivalent widths are being checked or newly determined. It is hoped that the manuscript will be ready for the general assembly of the International Astronomical Union at Moscow next year.

In describing this program, Dr. Minnaert discussed the particularly difficult problem of determining equivalent widths of the lines that are crowded together in the ultraviolet region. In fact, they overlap so much that the true continuous background of the solar spectrum can not be observed. The background can be inferred, however, from special study of the wings of strong lines of iron and other

elements, which differ markedly from the classical theory. Dr. Minnaert finds that in the region between wave lengths 3550 and 3650 angstroms the true background is about 40 per cent higher than the highest points observed.

Luminosities of Mira Stars

The long-period variables, of which Mira (α Ceti) is a well-known example, are giant red stars whose brightnesses change by five magnitudes, on the average, in cycles of the order of a year. Phillip C. Keenan, of Perkins Observatory, reported on his observations of the spectra of 20 such stars with the 60-inch reflector at Mount Wilson. His purpose was to determine their intrinsic luminosities, as indicated by the intensities of spectral lines that can serve as luminosity criteria.

A provisional calibration of the luminosity sequence shows that Mira variables of spectral class M2e, with an average period of 200 days, attain visual absolute magnitude -2.3 at maximum light. Later spectral types have lower luminosities and longer periods; thus, a typical M6e variable has a 360-day period and a visual absolute magnitude of 0.0.

Dr. Keenan finds that some long-period variables, such as RT Cygni and Z Ophiuchi, appear spectroscopically to be at least as luminous intrinsically as supergiants like Betelgeuse. The spectrum of Z Ophiuchi is noteworthy for the weakness of its calcium lines.

Nova Binary Star

Normally a 10th-magnitude star, the recurrent nova T Coronae Borealis has twice brightened briefly to the 2nd magnitude, in the years 1866 and 1946. It is also a close double star, the blue component responsible for the nova outbursts being accompanied by a giant red star of spectral type gM3.

Many new facts concerning this unusual system have been revealed in a spectroscopic study by Robert P. Kraft, Goethe Link Observatory, using the 100-inch Mount Wilson reflector. The period of binary revolution is 227.8 days. The motion of the red star was detected by R. F. Sanford a decade ago, but Dr. Kraft has now succeeded in finding spectral features of the blue star. Its hydrogen-beta line is bright, and shows a range in radial velocity of 70 kilometers per second.

From this it appears that the M-type star is about four times as massive as the sun, and the nova component has a mass of about 2.5 to 3 suns. The binary system is dynamically unstable, material flowing

from the red star to the blue one, forming a ring or shell around the latter. In addition, a large and very tenuous gaseous envelope surrounding the binary system as a whole is proposed by Dr. Kraft to account for the presence of certain spectrum lines of doubly ionized neon.

This model of the T Coronae Borealis system closely resembles present views of the nature of the variable stars SS Cygni and AE Aquarii, and Dr. Kraft believes that all three systems belong to the same homogeneous physical group.

Infrared Observations of Planets and Satellites

At McDonald Observatory, Gerard P. Kuiper has been continuing to observe bodies in the solar system with the 82-inch reflector in infrared light. He has measured the polarization of the light of Venus at wave lengths of one and two microns; the results appear to rule out water droplets as forming the planet's clouds (*Sky and Telescope*, November, 1954, page 20). Instead, Dr. Kuiper suggests the clouds of Venus may be made of particles of polymerized carbon suboxide (C_3O_2).

Striking differences among the four Galilean satellites of Jupiter have been




This astronomer from the Netherlands, M. G. J. Minnaert, was guest speaker at the society dinner at Urbana. Director of Utrecht Observatory, he is an authority on the sun's spectrum, eclipses, and stellar atmospheres. He is author of the popular book, "Light and Colour in the Open Air."



Another overseas visitor to the Urbana meeting was A. R. Hogg, who is a staff member of the Australian Commonwealth Observatory, on Mt. Stromlo near Canberra. He has been an active observer of southern double stars, but more recently his main field of interest has been photoelectric photometry.



Members and guests at the 98th meeting of the American Astronomical Society, University of Illinois, Urbana, August 18-21, 1957.



KEY TO PHOTOGRAPH

1 C, 2 L, 3 T, 4 R, 5 W, 6 J, 7 J, 8 W, 9 C, 10 H, 11 W, 12 W, 13 H, 14 H, 15 W, 16 R, 17 H, 18 S, 19 C, 20 W, 21 R, 22 S, 23 S, 24 R, 25 M, 26 R, 27 R, 28 W, 29 H, 30 C, 31 D, 32 H, 33 M, 34 K, 35 S, 36 H, 37 F, 38 G, 39 R, 40 A, 41 O, 42 E, 43 W, 44 H, 45 A, 46 F, 47 H, 48 W, 49 E, 50 H, 51 H, 52 S, 53 S, 54 K, 55 W, 56 M, 57 H, 58 M, 59 H.

60 P, 61 S, 62 F, 63 M, 64 W, 65 W, 66 B, 67 K, 68 M, 69 H, 70 C, 71 H, 72 W, 73 H, 74 K, 75 K, 76 S, 77 M, 78 D, 79 S, 80 M, 81 M, 82 A, 83 L, 84 S, 85 S, 86 M, 87 B, 88 K, 89 S.

90 B, 91 M, 92 S, 93 G, 94 U, 95 H, 96 S, 97 M, 98 S, 99 C, 100 V, 101 B, 102 S, 103 R, 104 K, 105 M, 106 L, 107 M, 108 S, 109 B, 110 F, 111 C, 112 M, 113 H, 114 B, 115 P, 116 Y, 117 S, 118 P, 119 S.

120 F, 121 M, 122 C, 123 R, 124 G, 125 B, 126 M, 127 E, 128 D, 129 M, 130 B, 131 H, 132 S, 133 M, 134 K, 135 K.

ALPHABETICAL LIST

40, H. Abt, Ye; 82, H. Albers, B; 45, L. Aller, M.

66, H. Bibcock, MWP; 114, J. Bartlett, Y; 90, W. Beardsley, A; 125, W. Bidelman, Li; 109, D. Billings, HA; 87, V. Blanco, CIT; 101, R. Bless, M; 130, D. Brouwer, Y; 111, D. Camp, I; 30, J. Caplan, Ad; 99, E. Carpenter, Steward Obs.; 70, J. M. Chamberlain, AMH; 122, J. W. Chamberlain, Ye; 19, G. Clemence, N; 1, Mrs. Clemence; 9, H. Crull, B.

128, D. Dewhirst, Cambridge Obs., England; 78, A. Dollfus, Paris Obs., France; 31, J. Douglas, Y.

127, F. Edmondson, I; 49, Mrs. Edmondson; 42, Miss M. Edmondson.

120, C. Federer, H; 110, J. Fernie, I; 37, R. Fleischer, RPI; 46, K. Franklin, AMH; 62, O. Franz, NW.

38, A. Gehrels, I; 93, Miss G. Gjellestad, Bergen, Norway; 124, R. Grenchik, V.

44, R. F. Haupt, N; 51, Mrs. Haupt; 71, R. W. Haupt; 131, J. Heard, DD; 13, C. Hemenway, Union; 14, Mrs. Hemenway; 10, D. Hemenway; 57, Miss S. Hemenway; 47, P. Herget, Cin; 36, Mrs. Herget; 113, A. Hewish, Cavendish Lab., Cambridge, England; 32, Miss S. Hill, Wellesley; 95, W. Hiltner, Ye; 17, A. Hoag, N; 59, A. Hogg, Mt. Stromlo Obs., Australia; 69, H. Horak, K; 29, C. Huffer, Wisconsin; 50, Mrs. Huffer; 73, J. Hynek, S.

7, A. Joy, MWP; 6, Mrs. Joy.

104, E. Kane, Detroit, Mich.; 88, Mrs. Kane; 34, G. Keller, Perkins Obs.; 75, I. King, Ill; 74, Mrs. King; 67, D. King; 54, R. Kraft, I; 134, J. Kraus, Ohio State; 135, G. Kuiper, Ye.

106, W. Liller, M; 83, P. Lind, Ad; 2, Miss J. Lucas, Ill.

129, T. Matthews, MWP; 121, Mrs. A. Matthews, MWP; 107, S. McCuskey, CIT; 63, D. McNamara, Brigham Young; 58, G. McVittie, Ill; 68, Mrs. McVittie; 105, T. Menon, H; 97, J. Merrill, Pennsylvania; 133, P. Merrill, MWP; 81, Mrs. P. Merrill; 25, M. Miller, S. J., Rochester, N. Y.; 33, W. Miller, S. J., Fordham; 56, R. Minkowski, MWP; 86, M. Minnaert, Utrecht, Holland; 91, W. Mitchell, M; 80, W. Moore, Louisville; 112, G. Moreton, Pomona, Calif.; 126, W. Morgan, Ye; 77, G. Mulders, ONR.

41, J. O'Keefe, Army Map Service.

60, J. Pawsey, Radiophysics Lab., Sydney, Australia; 118, B. Peery, M; 115, P. Pesch, Ye.

27, E. Rabe, Cin; 21, Mrs. Rabe; 26, Miss I. Rabe; 4, Miss K. Rabe; 39, R. Redlich, RPI; 103, R. Riddle, Purdue; 24, Miss M. Risley, Randolph-Macon; 123, W. Roberts, HA; 16, M. Rogers, Ill.

108, J. Schopp, Missouri; 102, Mrs. Schopp; 119, L. Searle, DD; 117, C. Seyfert, V; 92, A. Shatzel, Ad; 85, B. Sitterly, American Univ.; 96, Mrs. C. Sitterly, Nat'l. Bur. of Standards; 35, H. Smith, Y; 132, A. Spitz, S; 79, R. Sprague, K; 53, J. Stebbins, Li; 76, J. Stokley, Michigan State; 89, K. Strand, NW; 84, Mrs. Strand; 61, Miss C. Strand; 18, Miss J. Streeter, ONR; 22, B. Stromgren, Ye; 98, O. Struve, Le; 52, C. Swanson, Army Ballistic Missile Agency, Huntsville, Ala.; 23, G. Swenson, Ill.

3, S. Trehan, Ye.

94, A. Uppgren, M.

100, M. Vardya, M.

28, N. Wagman, A; 11, Mrs. Wagman; 55, H. Weaver, Le; 72, L. White, ONR; 20, R. Whitehurst, Alabama; 43, G. Wilkins, N; 5, Mrs. Wilkins; 8, M. Wilkins; 64, R. Wilson, Naval Res. Lab.; 65, Mrs. Wilson; 48, Miss K. Wilson; 12, S. Wyatt, Ill; 15, Mrs. Wyatt.

116, K. Yoss, Louisiana State.

A, Allegheny; Ad, Adler Planetarium; AMH, American Museum-Hayden Planetarium; B, Butler; Cin, Cincinnati; CIT, Case Institute of Technology; DD, David Dunlap; H, Harvard; HA, High Altitude; I, Indiana; Ill, Illinois; K, Kansas; Le, Leuschner; Li, Lick; M, Michigan; MWP, Mount Wilson-Palomar; N, Naval Observatory; NW, Northwestern; ONR, Office of Naval Research; RPI, Rensselaer Polytechnic Institute; S, Smithsonian Astrophysical; V, Vanderbilt; Y, Yale; Ye, Yerkes.


found from intensity tracings of their spectra with an infrared spectrometer. Satellites I and IV reflect sunlight with little change, but for III and especially II the spectrum is markedly fainter in the range from 1.5 to 2.5 microns. This effect would be produced if the surfaces of satellites II and III were covered with snow. The high reflectivity of II in visual light and its color fit this hypothesis, while III, which is darker, may be covered with snow contaminated by silicate dust. At Jupiter's distance from the sun, the rate of evaporation of snow would be very slow, even if exposed to a vacuum.

In infrared spectrum tracings of Mars, Dr. Kuiper has observed three absorption bands of carbon dioxide near two microns, finding them stronger than can be explained by the amount of this gas in the earth's atmosphere. This fact was established by a comparison of the tracings reproduced here, which were made when the light of Mars and the sun passed through nearly equal thicknesses of our air. At the left each tracing shows the same deep absorption band at 1.8 microns produced by water vapor in the earth's atmosphere. The three CO_2 absorptions are ω_0 , ω_1 , and ω_2 , at wave lengths 1.957, 2.006, and 2.057 microns, respectively. Relative to the sun's spectrum, the Martian bands are stronger, in approximately the same ratio as were weaker bands of carbon dioxide at 1.6 microns discovered by Dr. Kuiper in 1947. Thus the new results corroborate the earlier evidence for CO_2 as a constituent of the Martian atmosphere.

Other McDonald observations confirm that Saturn's rings are composed of snow particles, and reveal new features in the infrared photographic spectra of Jupiter and Uranus.

Brighter Artificial Satellites?

Because a 20-inch American artificial satellite is expected to be about 6th magnitude at perigee and about 10th at apogee, its faintness will be an important difficulty in optical observations. Armand N. Spitz, of the Smithsonian Astrophysical Observatory, and Raymond H. Wilson, Jr., Naval Research Laboratory, propose



These spectrum tracings by G. P. Kuiper show the added strength of carbon dioxide bands in the spectrum of Mars. Yerkes-McDonald Observatories chart.

that the satellite be a polyhedron, with plane mirror surfaces.

Illuminated by the sun, such a satellite would be visible as a series of light flashes three or more magnitudes brighter than the reflection from a sphere. Dr. Wilson has computed that a plane mirror surface only one centimeter in area, situated about 300 miles from the observer and reflecting sunlight face-on, could appear as bright as a 1st-magnitude star.

The rate of occurrence of the flashes would depend on the number of plane faces and the speed of the satellite's rotation. In an alternative design, plane facets would be added to the original sphere, so the satellite would remain visible during the intervals between flashes, which would occur with a frequency of several times a second to several times a minute.

Streamlining of the satellite's surface is not necessary, because of the low air resistance at the heights where it will travel in its orbit. While being carried upward from the earth, the satellite will be enclosed in the third stage of the Vanguard rocket, so during that time also its shape will not affect drag.

Both speakers pointed out that a flashing body might be easier to recognize as a satellite by visual observers, and the flashes could tell how fast it was rotating.

Punch-Card Observing

At the Flagstaff, Arizona, station of the U. S. Naval Observatory, an observer at the 40-inch reflector is assisted by automatic devices that punch the data of each photoelectric observation directly onto IBM cards. These can then be sent to headquarters in Washington, D. C., for processing by IBM 650 computing machines.

Two separate analyzer sections of the photometer may be readily interchanged, one being used for observations of star brightnesses in three selected colors, the other for polarization measurements. The photometer output is fed through amplifiers to an integrator, and the result is displayed on a strip-chart recorder.

But the recorder is also equipped with a digitizer and readout device which punches on IBM cards the amount of each deflection of the recording pen, with an accuracy of a thousandth of the full-scale deflection. Observations of a particular star are made automatically in sequence after the telescope is set by the observer. Fifteen to 20 stars can be observed in an hour because the telescope is so easily pointed from one part of the sky to another. (See *Sky and Telescope*, November, 1956, page 4.)

Interstellar Gas

Sidney van den Bergh, of Perkins Observatory, calls attention to a surprising property shared by both the Milky Way galaxy and the Andromeda nebula: the relatively small amount of interstellar gas in the spiral arms. While ejection of matter from stars provides some such gas, a larger quantity is consumed in the process of star production. Were no external supply available, the gas in the solar vicinity would be exhausted in about 7×10^8 years.

Furthermore, the amount of gas now in the galactic nucleus is only about 1/250 or 1/300 that which should have been ejected from the stars there during the five-billion-year history of our galaxy.

Dr. van den Bergh suggests that both these points can be explained by a continuing escape of gas from the nucleus, the gas moving outward along the spiral arms. The present rate of gas loss from the central part of the galaxy would match the rate at which the interstellar gas is being removed from the spiral arms by star formation.

The Seeing at Kitt Peak

A. B. Meinel, of the new National Astronomical Observatory, has encouraging data on the quality of the seeing on Kitt Peak, Arizona, one of the five sites in the Southwest being tested for the location of 80-inch and 36-inch reflectors (*Sky and Telescope*, August, page 482).

The seeing on Kitt Peak, an isolated mountain mass rising to 6,875 feet, with



The present director of the University of Illinois Observatory, George C. McVittie, stands at the left in this picture. Talking with him are two of his predecessors in office, Joel Stebbins (center), who was director from 1903 to 1922, and Robert H. Baker, from 1922 to 1951. Photograph by Walter J. Miller, S. J.


a 70-acre level summit, has proved of surprising excellence. This was shown by the small amount of scintillation "noise" on the photoelectric records of Polaris, which was continually observed with a 6-inch monitor telescope. The Kitt Peak observers believe that the seeing is better there than at any other observatory where they have worked. Clouds are rare, especially in the summer, when there is less cloudiness than at nearby Tucson. Another advantage is the smallness of the day-to-night temperature range — only 3.1 degrees centigrade in winter and 6.5 degrees in summer.

Thirty tons of equipment and a housing trailer have been hauled to the summit of Kitt Peak over a primitive bulldozer trail which has grades as steep as 78 per cent. In July, a 16-inch Cassegrainian reflector, equatorially mounted on a truck, was put into service. The seeing tests at Kitt Peak and at the other stations under survey will continue well into the future.

Expanding Star Cluster in Orion

The compact cluster of faint stars surrounding the Trapezium, in the heart of the Orion nebula, is growing in size at the rate of one part in 300,000 per year, according to the work of K. A. Strand, director of the Dearborn Observatory, Northwestern University. This expansion indicates a common origin for the cluster stars only 300,000 years ago, with an uncertainty of 40,000 years.

Dr. Strand's findings illustrate the importance of positional observations of the stars over long intervals. He made measurements of recent Orion photographs taken with the 40-inch Yerkes refractor, and compared them with similar plates



This map of the surroundings of the region of the Trapezium (Theta Orionis) includes stars to visual magnitude 17.5. Dr. Strand based the chart on infrared photographs taken at Lick Observatory and the Flagstaff station of the U. S. Naval Observatory. The coordinates are for the epoch 1900.

taken 50 years earlier by G. W. Ritchey and J. A. Parkhurst. Since all the exposures were made through yellow filters on yellow-sensitive plates, the nebulosity was so weakened that it did not interfere with the measuring of the star images.

From 10 pairs of plates with 50-year intervals, precise proper motions were determined for 200 stars of the 14th visual magnitude and brighter, within an area roughly half a degree square. These measurements were made with respect to 16 reference stars of average magnitude 11 that belonged to the cluster. The configuration of the 16 stars turned out to be one part in 6,000 larger on the recent plates than on the early ones.

Was this change due to an actual radial expansion of the cluster? One possible alternative was that the apparent expansion was merely the result of random motions of the reference stars. However, new measurements with another set of 25 reference stars showed the same enlargement. Other tests demonstrated that there was no instrumental effect, such as a progressive increase in the focal length of the Yerkes telescope, nor was there a gradual shrinkage of the emulsion on the old plates. When Dr. Strand compared two pairs of photographs of the Pleiades taken 50 years apart with this same telescope, no expansion was detected.

By comparing his new very precise proper motions of the cluster stars with previously published radial velocity values, Dr. Strand computed 600 parsecs as the distance of the cluster. This is considerably greater than the 400-parsec result derived photometrically by P. P. Parenago in Russia, and by W. A. Hiltner and H. L. Johnson in the United States. But Dr. Strand believes the latter determinations

are too small because the A-type stars in the Orion cluster are actually more luminous than their colors imply.

Within five minutes of arc of the Trapezium there are seen about 300 stars brighter than visual magnitude 17.5. When allowance is made for stars hidden by patches of interstellar obscuring material, the number becomes about 400, corresponding to a total mass of some 350 suns — a minimum value. This is nearly the same as the mass of the Pleiades cluster, but only within one-eighth of the volume.

With an age of only 300,000 years, the Trapezium cluster is among the very youngest known. This is confirmed by the color-magnitude diagram of its members constructed by Dr. Strand, showing to an exaggerated degree the peculiarities found by Merle Walker for the young cluster NGC 2264 (*Sky and Telescope*, October, 1954, page 426), which is now believed to be three million years old.

The color-magnitude diagram of the Trapezium cluster has a broad, continuous sequence of stars from spectral type O6 to type K5 V. But the stars later than B3 are brighter than standard main-sequence stars, showing that they may still be contracting gravitationally.

QUESTIONS... FROM THE S+T MAILBAG

Q. To what distance does the absolute magnitude of a star correspond?

A. The absolute magnitude of a star is the magnitude it would appear to have if viewed from a standard distance of 10 parsecs (32.6 light-years).

Q. When can the constellation Crux (the Southern Cross) be seen from southern Florida?

A. It is best placed for view when the quadrilateral of Corvus is on the meridian, Crux then being 40° to the south. This can happen during hours of darkness from January until June, specifically at 11 p.m. local time at the beginning of April, and 9 p.m. at the beginning of May.


Q. What is meant by the radial velocity of a star and why can it have positive and negative values?

A. Radial velocity is the speed at which a star approaches us (negative value) or recedes (positive value); it is that part of the star's motion that lies along our line of sight to the star.

Q. How is radial velocity measured by astronomers?

A. By the Doppler shift in the positions (wave lengths) of lines in the spectrum of the astronomical body. Motion of approach shifts the spectral lines to the violet side of their normal positions, recession shifts them toward the red. The amount by which a spectral line is displaced is directly proportional to the radial velocity.

W. E. S.



Note the predominance of outward motions in this Dearborn Observatory plot, in which the circles centered on the Trapezium have radii of five and 10 minutes of arc, respectively. For the number of stars indicated by the adjoining label, each arrow gives the average proper motion; the arrow in the box has the unit length of one second of arc in 1,000 years.

A Seeing Compensator Employing Television Techniques

JOHN H. DEWITT, ROBERT H. HARDIE, and CARL K. SEYFERT, *Dyer Observatory, Vanderbilt University*

THE BASIC PROBLEM in photographing the surface of a planet is overcoming the disturbances in the earth's atmosphere. It has always been possible to see more detail visually than can be recorded photographically through the same telescope. This is due to the fact that the eye can distinguish between moments of atmospheric steadiness and other instants when the image is blurred.

There has long been a pressing need for a photographic method that can record the fine planetary detail revealed to experienced visual observers in instants of excellent seeing. This could be accomplished by making extremely short, high-contrast exposures with a moderate-size telescope. Instruments of up to 24 inches are more likely to produce good images during times of atmospheric unsteadiness than are larger telescopes. Since smaller apertures ordinarily require longer exposures, some method of image intensifica-

tion is required. The apparatus described here, which uses a closed-circuit television system on a 24-inch reflector, not only achieves the desired image intensification and added contrast but compensates electronically for the movement of the planetary image due to seeing.

The Orthicon pickup tube used in television cameras is a very powerful light amplifier, if the illumination on the photocathode is sufficient to overcome beam noise. This condition is satisfactorily obtained with the Dyer 24-inch reflector on the planets Venus and Jupiter, and also Mars when it is near opposition.

The telescope is used in its Cassegrainian form, and light passes through a negative lens to form an image of Jupiter about half an inch in diameter. As the block diagram shows, the beam of light is divided by a dichroic filter so that about 60 per cent is fed to the photocathode of the Orthicon, while the remainder forms

a monitor image on two slits placed at right angles to each other.

The planetary image on the photocathode produces on the monitor Kinescope a picture approximately four times larger and much brighter than the original. The modified TV circuit is so arranged that the system can be turned on at intervals of 1/30 second, the time corresponding to one vertical sweep. Other circuits allow the Orthicon target to store for 1/15, 2/15, or 4/15 second before the single-sweep discharge is applied. Such accumulation of electrons by storage on the target greatly increases the sensitivity to faint light.

A primary purpose of the new device is to compensate for the movements of the image as a whole during the exposure. To achieve this, four deflection coils are placed around the image section of the Orthicon at 90-degree intervals. Currents of the proper phase and strength pass



The seeing compensator is the T-shaped device mounted at the Cassegrainian focus of the 24-inch reflector. In the cross of the T is a beam splitter that diverts part of the planet's light down the side arm to the Orthicon television pickup tube. The remaining light continues along the top of the T to crossed slits and two photocells, which detect image shifts and provide for their compensation. Cables run from the telescope assembly to a cabinet for the power supply and other electronic components. The light-colored disk on the cabinet is the Kinescope screen, where the intensified image is displayed. Photos by Jeanne-Gordon Studio.

through these coils, and deflect the electron stream between the photocathode and target in such a way that the image remains centered on one spot on the target, even though the optical image on the photocathode may wander across 1/10 the diameter of the field of view. The currents that actuate the deflection coils come from two 1P21 photocells, which are placed behind the two slits at the second or monitor image of the planet.

These photocurrents are directly proportional to the image position in x and y co-ordinates along each slit. The currents pass through d.c. amplifiers which use vacuum tubes and high-power transistors, and then are fed to the image-section deflection coils. The coils must be aligned carefully in order to have their action just offset the wandering of the optical image. This is done by adjusting the deflection-compensator amplifier gains so that the image is stationary. One simple way is to shake the telescope and adjust the gains until the image movement is compensated!

The seeing compensator, or "tranquillizer" as it has been called, is pictured here, as mounted on the 24-inch reflector. The cabinet holds the power supplies, synchronizing generator, amplifiers, and monitor Kinescope. To photograph the image we use a 35-millimeter camera (not shown), which is mounted about nine inches in front of the television screen.

It has been a relatively short time since the apparatus became ready for operation, and up to September its principal use was to take pictures of Jupiter, samples of which are reproduced here. The images on this page were taken through the apparatus with the compensator operating (left), and with it turned off (right).


A television circuit of this type can

utilize the full defining power of the telescope, because it employs more scanning lines than does present-day commercial TV. In our apparatus, the image of a planet contains 200 scanning lines which, in the case of Jupiter, are spaced 0.2 second of arc apart—about the theoretical

period; also we expect to photograph Mars at its next opposition.

Another project that appears feasible is to test the coherence of seeing—the tendency of two separated stars to be simultaneously affected the same way by seeing. The test could be made by steadying the

Photographs of Jupiter on the Kinescope screen, with the shadow of one of its larger moons seen against the planet's disk. The pictures at the right were taken with the seeing compensator turned off, those at the left with it in operation. These examples do not fully exploit the definition of the apparatus, but they demonstrate the steadying effect of the compensator. Dyer Observatory photographs.



resolving power for a 24-inch telescope. It seems possible to see readily on the TV screen as much as an experienced observer does at the telescope eyepiece. Increased contrast can be obtained through the electronic system, rendering faint detail much more apparent than is the case with direct telescopic viewing.

In addition to photographing Jupiter, the apparatus has been used successfully as an automatic guiding device. This fall and winter it is planned to observe Venus, in an attempt to evaluate its rotation

image of Jupiter and examining the fluctuations of its satellites at various angular distances from the planet.

Our apparatus was developed as a joint project of Vanderbilt University and Station WSM-TV in Nashville, Tennessee.


JUPITER'S CLOUDS

It is widely agreed by astronomers that the atmosphere of Jupiter above the cloud layer that veils the planet consists mainly of hydrogen and helium, with small quantities of methane and ammonia. The clouds themselves probably are made up of particles of frozen ammonia.

P. Squires, in Australia, now finds that the equatorial clouds of Jupiter are not flat-topped, as had been previously assumed, but are domed like terrestrial cumulus clouds. The cumuliform tops reach about 20 or 30 kilometers above the main cloud deck. Dr. Squires reached these conclusions from a theoretical analysis of observations, made several years ago by Seymour Hess, of the intensity of spectrum bands of methane and ammonia over different portions of the disk of Jupiter.

The new study indicates that the temperature at the level of the main deck is considerably higher than -150° Fahrenheit. However, with the introduction of this new and more complicated model of Jupiter's clouds, it is no longer possible to deduce an accurate temperature for the cloud surface from the spectroscopically measured quantity of ammonia.

Dr. Squires, who is a staff member of the division of radiophysics, Commonwealth Scientific and Industrial Research Organization, Sydney, Australia, reported his findings in the *Astrophysical Journal*.



The operation of the seeing compensator is indicated in this schematic diagram of the optical and electronic components.



A Probable Meteorite Fall in Brazil

VINCENT MENEZES

This fragment of the meteorite was found near Ibitira. Its rough dimensions are 4 by 6 or 12 inches. The dark, shiny surface is part of the crust; the lighter portion at the left is where a sample was taken for analysis. Photo by Geraldo C. Lessa.

ON the late afternoon of June 30, 1957, between 5:00 and 5:30 p.m. local time, a great fireball passed northwestward over the central part of the state of Minas Gerais, Brazil, and sounded as if it were exploding in the air, frightening nearby villagers.

As soon as our amateur group, the Center for Astronomical Studies of Minas Gerais, learned of the event, we appealed through the local press and by letter for observers' reports. Of the 32 letters received, we selected the best 19 and made field trips with notebooks, maps, and compasses, to interview the writers and others who had seen the phenomenon.

Observers in this city, Belo Horizonte, first saw the fireball at an altitude of 50 degrees; falling toward Martinho Campos and seeming to explode in the air. Other persons, about 160 kilometers to the southeast of the end point of the fireball, saw the breakup take place about five degrees above the horizon. From esti-

mates of this kind it appears that the visible trajectory came to an end about 10 or 12 kilometers above the earth's surface.

The fireball was seen as a reddish, egg-shaped body, which became silvery in color, but I think the latter hue was seen after the disintegration. There was a noise like the reverberation of thunder. The accompanying sketches show the long white train which the meteor left behind it, as it appeared at first and after it had become distorted by upper-air winds. The train seemed straight to observers directly beneath it, and as a parabolic arc to those who viewed it from the side. The straight train became sinuous, and then divided in two, taking on a V-like shape, its color becoming silvery blue and light gray.

Several kinds of evidence fixed the end point of the trajectory in the county of Martinho Campos, near Ibitira, a village of some 200 inhabitants, on the Minas Gerais state railroad 213 kilometers from Belo Horizonte. The directions in which



the trail was seen by different observers are shown by arrows on the map, and they tend to converge on Ibitira.

At the Monjolo farm near Ibitira, the characteristic thunder sound was heard, followed some seconds later by a whistling like a falling bomb, and immediately thereafter by the noise of something like a stone striking the ground. A young man on the farm looked at once toward the direction from which this sound seemed to come, and saw cattle running down a hillside pasture into the valley.

At first no meteoritic fragment could be located, for there were too few people to scour the many acres of pasture, which are covered by creeping vegetation. Later, however, the small piece shown in the picture was found in Ibitira near the point of the fall predicted by my analysis. It lay in a hole in the ground about 10 inches deep and eight in diameter.

This object has a brilliant black crust, and a light brown, slightly porous interior. It was brought to the Institute of Technological Researches, Belo Horizonte, for chemical analysis. Preliminary results show the presence, in decreasing order of abundance, of silicon, magnesium, iron, aluminum, calcium, chromium, and titanium. Spectroscopic examination reveals lines of sulphur and manganese. Contrary to expectation, there were no signs of sodium, nickel, or cobalt.

These facts make it probable that this object is actually a stony meteorite. A further search is intended to recover the fragment that we suppose must have fallen on the Monjolo farm.




Above: The upper row of sketches shows the original appearance of the meteor train, and the lower row the effect of wind distortion soon afterward, for observers looking toward: I, the northwest; II, the northeast; III, the southwest; and IV, from Ibitira.

Left: Observers' reports of the direction of the end point of the meteor's trail are indicated by the arrows on this map. At Ibitira, the meteor seemed to be coming directly toward the observer.

ARTIFICIAL SATELLITE NO. I

These are photographs of the screen of an oscilloscope fed by a short-wave receiver, showing 20,005-megacycle pulses from the Russian satellite, during the night of October 5, 1957, one day after the launching. The various pulse patterns may indicate actual telemetering, or merely effects of propagation through the atmosphere. Photos by Robert Slavin and G. R. Miczaika, Geophysics Research Directorate, Air Force Cambridge Research Center, Bedford, Massachusetts.



THE ARTIFICIAL-SATELLITE AGE began on October 4, 1957, when a Russian test vehicle started to circle the earth 15 times a day.

According to a preliminary analysis by German astronomers at Bonn Observatory using early observations, the new moonlet was launched that day about 20^h Universal time from a point somewhere in central Asia.

In *Soviet Aviation*, Prof. Y. A. Pobedonostsev stated that the three-stage launching rocket attained a speed of about 4,500 miles per hour in the first two minutes. When the first stage dropped away, the vehicle was moving upward at a 45-degree angle. At second-stage burnout, the speed was nearly 12,000 miles an hour. Unpowered flight lasted until horizontal motion was achieved about 625 miles from the launching site; then the third-stage motor took over to boost the speed to approximately 18,000 miles per hour before ejection of the artificial moonlet.

During the first days of the satellite, conflicting reports and speculation were mingled with facts, and the confusion has been only partly cleared away at this writing. But we know, for instance, that the satellite is a sphere 22.8 inches in diameter, with the surprisingly large total weight of 184 pounds. According to Radio Moscow on October 5th, the period of orbital revolution was 96.2 minutes, implying an average height of 370 miles.

As soon as the existence of the moonlet was revealed, MOONWATCH teams were alerted; then it was learned that no favorable passages over the United States would occur for a number of days. Like the projected American satellites, the Soviet one presumably can be seen only during or near morning and evening twilight, when it may be viewed as a sunlit speck against a fairly dark sky; it would be invisible at other times.

American observations, at least initially, were limited to the radio signals transmitted from the satellite, the continuous beeping audible over a range of thousands of miles because of the great altitude of

the satellite. The transmitters, which consumed one watt of power, radiated pulses of 0.3-second duration, followed by an equal interval of silence, at 20,005 and 40,002 megacycles (wave lengths of about 15 and 7.5 meters). The pulses at one frequency occurred simultaneously with the pauses on the other. By October 8th, these transmissions appeared to be growing weaker.

The great early strength of the signals made them easily detectable by amateur operators, and they were received by many members of the American Radio Relay League. The 10 Minitrack stations organized by NRL to monitor American satellites at 108 megacycles required hasty conversion of antennas and receivers to pick up the Russian frequencies.

During a passage of the satellite, the steady beeping could be followed for 25 to 35 minutes. At one crossing over the New York City area, at 7:52 a.m. EST on October 5th, the receiving station of RCA Communications, Inc., at Riverhead, Long Island, measured the change in frequency of the carrier wave as the satellite approached and then receded, an example of the Doppler effect. From this the orbital velocity was deduced to be 4.92 miles per second.

Several monitoring stations picked up other strong 20-megacycle signals, probably originating from a ground station in the Moscow area, which apparently triggered the telemetering system in the satellite.

The earliest reports of visual sightings were contradictory. Later, apparently reliable optical observations of a bright object were obtained on October 6th and 8th at Mt. Stromlo Observatory, Australia, and at the University of Alaska. On the morning of October 9th, it was seen as a 5th-magnitude object from the University Observatory, Vienna, Austria.

Smithsonian astronomers obtained a preliminary orbit on October 9th, using the digital computer at Massachusetts Institute of Technology. Their calculations, based on a radio observation at Boulder,

Colorado, and sightings in Alaska and Australia, gave 370 miles as the average height of the satellite. Later computations gave a perigee distance of 143 miles and an apogee of 583 miles.

Actually, three pieces of "hardware" were moving in closely similar orbits: the satellite itself, the discarded third stage, and the empty nose cone of the rocket. Probably most authenticated visual observations refer to the third stage which, much larger than the 22.8-inch sphere, may well appear brighter. Because of its greater air drag, the third stage should be moving below and *ahead* of the satellite, a seeming paradox predicted by celestial mechanics. This fits a report from Hobart, Tasmania, of the passage of a bright body followed by a fainter one.

The first indications of the downward spiraling and eventual destruction of the Russian moonlet may have been reports from the Cavendish Laboratory, Cambridge, England, that radio observations showed a diminished height. At this writing, the lifetime of the satellite cannot be foretold with accuracy.

The first MOONWATCH observation seems to be that by James Plato, at New Haven, Connecticut, on October 10th at 5:23 a.m. EST, confirmed by station leader Robert Brown. The satellite (or more probably the third stage) was photographed on October 9th with the super-Schmidt meteor camera at Newbrook, Alberta, Canada. On October 12th, the Amateur Telescope Makers of Boston timed an early-morning passage of the third stage across the fields of several telescopes, getting a good taped recording.

Observers with MOONWATCH equipment should be especially on the lookout for the fainter true satellite, of probably 5th to 7th magnitude, which was trailing the sunlit third stage by many minutes.

Other satellites, both American and Russian, are expected during the remainder of the International Geophysical Year. The first American launching is to be in December, according to President Eisenhower's statement of October 9th.

NEWS NOTES

SATELLITE WEATHER PICTURES TO BE MADE WITH PHOTOCELLS

Hurricane warnings and other vital weather information could be more comprehensive and accurate if we could obtain views of the earth from artificial satellites. This has already been proposed by Harry Wexler (*Sky and Telescope*, March, 1956, page 208). Last spring he released a picture showing the hypothetical appearance of North America and its cloud formations from a television camera in a satellite 4,000 miles high.

Such a television "station" in space is beyond present technological achievement, but in August's *Scientific Monthly* S. F. Singer, University of Maryland, describes how simple photocells could be used to scan the brightness changes of clouds, continents, and oceans, from a satellite only 200 or 300 miles above the earth's surface.

Repeated transmissions from the satellite photocells to ground stations would permit the building up of weather pictures in which trained meteorologists could recognize cloud formations and track their movements. Although this method would take longer than a television camera to record the picture of an area, weather movements are slow enough so this would not be a handicap.

The resolution required to recognize weather features need not be great — 10 to 100 miles would do. On the basis of Dr. Wexler's original picture, Dr. Singer shows how our continent would look if scanned, from a pole-to-pole orbit 300 miles high, by a single photocell making adjacent sweeps in a nearly north-south direction. The photocell would have a field of view of 800 kilometers at right angles to the satellite's motion.

In such a "photograph," large cloud masses could be recognized, while the same view taken with two photocells mounted in the satellite would show additional details. A four-cell "exposure" would clearly reveal the most important features of the television picture, including cyclones, West Indian hurricanes, and line squalls, with smaller details being moderately visible.

1959 TOTAL ECLIPSE

The next total eclipse of the sun to be visible from the United States will occur two years from now, on October 2, 1959. According to the detailed predictions in Circular 78 of the U. S. Naval Observatory, totality may be seen shortly after sunrise in northeastern Massachusetts and southern New Hampshire, in a strip about 38 miles from north to south and about 63 miles from east to west. The duration of totality will be about 56 seconds along the central line, which passes nearly through Fitchburg and Salem, Massachusetts. Boston lies inside the path of totality. In this

entire area, however, the sun will be less than two degrees above the horizon.

After leaving the New England coast, the path of totality spans the Atlantic Ocean, crosses North Africa, and ends in the Indian Ocean. The closing partial stages will be visible from a large part of the eastern United States, where the sun will rise already eclipsed.

ONR ASTRONOMY PROGRAM

The Office of Naval Research will continue its program in support of astronomical research during the year June, 1958, to June, 1959. Dr. G. C. McVittie, University of Illinois Observatory, is chairman of its seven-man advisory panel. Applications for support of research in astronomy must be submitted by December 15th. Details can be obtained from the Chief of Naval Research, Department of the Navy, Attn: Code 410, Washington 25, D. C.

DISTANCES OF CLUSTERS

New values for the distances of 16 prominent galactic star clusters are presented by Harold L. Johnson, Lowell Observatory, in the July issue of the *Astrophysical Journal*.

These distances are photometric, being based on photoelectric measurements of cluster stars in ultraviolet, blue, and yellow light. Through comparison of the three brightnesses for each star, he could determine both the amount of dimming by interstellar dust and the intrinsic color (hence absolute magnitude, in the case of a main-sequence star). Dr. Johnson has refined the method to take into account evolutionary changes in the brightnesses of stars of different ages. The distances found in this way have, for the most part, uncertainties between 10 and 20 per cent.

Dr. Johnson has also deduced photometric distances of four stellar associations: I Persei, 520 light-years; II Persei, 1,140; I Orionis, 1,300; and III Cephei, 2,380.

DISTANCES OF CLUSTERS

Name	Constellation	Parsecs	Light-years
Hyades	Taurus	40	130
Coma cluster	Coma Berenices	80	260
Pleiades	Taurus	126	410
Praesepe	Cancer	158	515
Messier 39	Cygnus	250	820
IC 4665	Ophiuchus	330	1,080
Messier 34	Perseus	440	1,440
Messier 25	Sagittarius	550	1,790
Messier 67	Cancer	830	2,710
NGC 2264	Monoceros	870	2,840
Messier 36	Auriga	1,260	4,110
NGC 2362	Canis Major	1,450	4,730
NGC 6530	Sagittarius	1,580	5,150
NGC 2244	Monoceros	1,660	5,410
Messier 11	Scutum	1,740	5,670
Double cluster	Perseus	2,250	7,340

IN THE CURRENT JOURNALS

THE ORIGIN AND IMPLICATIONS OF THE COSMIC RADIATION, by Serge A. Korff, *American Scientist*, September, 1957. "We may also say a word about the astrophysical implications contained in the galactic acceleration picture of cosmic rays. There are two quite different possibilities, first that cosmic rays are in equilibrium today, being produced, accelerated, and absorbed, so that we are measuring a part of a long-term equilibrium process, and the second that cosmic rays are residual from an original catastrophic explosion."

AIR NAVIGATION SINCE WORLD WAR II, by John F. Heard, *Journal of the Royal Astronomical Society of Canada*, August, 1957. "Since the end of the war, I believe few astronomers have had opportunities to keep abreast of the developments in air navigation. These developments have been rather astonishing, both from the point of view of principle and technique, and a review of them should be of interest to astronomers."

WEST VIRGINIA CEREMONY FOR RADIO OBSERVATORY



On October 17th at Green Bank, West Virginia, a ground-breaking ceremony was scheduled for the new National Radio Astronomy Observatory. This major research facility is to have a radio telescope 140 feet in diameter, as described on page 398 of the July, 1956, *Sky and Telescope*. The observatory is being sponsored financially by the National Science Foundation, which has made a contract for its establishment and operation with Associated Universities, Inc.

RADIO TEST FOR A LUNAR ATMOSPHERE

A new demonstration, more far-reaching than any before, of the extremely low density of any atmosphere the moon may have is reported in the August *Philosophical Magazine* by B. Elsmore, of the Cavendish Laboratory, Cambridge, England.

On January 24, 1956, the moon occulted the Crab nebula in Taurus, one of the most intense localized radio sources in the sky. This phenomenon was recorded with the Cambridge radio telescope on a wave length of 3.7 meters. As the diameter of the source at this wave length is about five minutes of arc, the observed radio intensity took some 10 minutes to decrease to zero as the moon moved in front of the nebula, and a corresponding increase took place gradually at immersion an hour later.

The total duration of the occultation was observed to be 59.6 minutes, with an uncertainty of ± 0.26 minute, as compared with the predicted duration of 59.2 min-



Sudden brilliant outbursts on the sun known as flares are key phenomena in the study of solar-terrestrial relations. These Sacramento Peak Observatory pictures, in hydrogen light, show a strong flare that occurred on September 18th. At 17:37 UT it was of importance 2+ (left), and by 18:38 had grown to importance 3+ (right). There were also sharp bursts of solar radio emission at 18 megacycles. The dark filaments are ordinary hydrogen prominences seen against the sun's disk.

utes. While the excess of 0.4 minute may not be significant, it could be interpreted as due to a refraction of the radio waves by free electrons in a lunar atmosphere. The electron density at the moon's surface required to produce the effect would be about 1,000 per cubic centimeter.

Dr. Elsmore points out that on the sunlit side of the moon whatever atmosphere there is should be completely ionized, all of its atoms broken up into ions and free electrons. Hence he could calculate the total amount of atmosphere necessary to provide an electron density of the above amount. It turns out that any permanent lunar atmosphere must have a density less than 5×10^{-13} that of the earth's air at sea level. This is a maximum value, since the observed amount of refraction of radio waves by the moon is an upper limit.

This radio result extends the work of Audouin Dollfus, who photographed the moon's cusps at the Pic du Midi Observatory and concluded that the lunar atmosphere must be less than 10^{-9} the density of air at sea level (*Sky and Telescope*, December, 1956, page 72).

LONG-ENDURING METEOR TRAINS

According to Charles P. Olivier, retired director of the Flower and Cook Observatory, one meteor out of each 750 leaves behind it a luminous train lasting for more than 10 seconds, as seen with the unaided eye. He has recently published a catalogue of 575 such long-enduring trains, supplementing his two earlier lists of 1,492 similar objects.

Of the 575 trains, observed chiefly between the years 1926 and 1956, six persisted for one hour or longer. In many cases the drift of the train across the sky allowed the determination of upper-air

Compare the sun's disk, as seen here in white light, with the hydrogen pictures above to find the relation of the flare area and the largest group of sunspots visible that day. This photograph was taken through clouds at the U.S. Naval Observatory, Washington, D. C., by Mrs. Winifred S. Cameron.



wind velocities, averaging about 200 kilometers per hour. The relatively few determinations of these winds from day and twilight meteor trains give slightly lower values.

Vivid colors are occasionally observed for the nighttime trains, but by day or in twilight the hue is ordinarily smoky or grayish. In recent years it has become increasingly difficult to distinguish between daytime trains and the much commoner vapor trails left by aircraft.

In the same article, Dr. Olivier presents a table of heights and orbital elements for 102 fireballs, as deduced from visual observations. While Baker-Schmidt cameras and radars give far greater accuracy, stations with such equipment are so few that only in rare cases can they secure observations of the most brilliant and interesting fireballs. Visual records by amateurs remain very desirable, especially for objects appearing over the oceans, which otherwise would not be reported.

Dr. Olivier's study appeared in the June, 1957, *Proceedings of the American Philosophical Society*.

NSF RESEARCH GRANTS

Proposals for funds to support professional research in astronomy may now be submitted to the National Science Foundation; if received by the end of November they will be considered at the annual meeting of the foundation's advisory panel.

A booklet giving detailed information on the preparation of proposals may be obtained from Dr. Geoffrey Keller, program director for astronomy, at the National Science Foundation, Washington 25, D. C.

SOUTH-NORTH COMPARISON

There are approximately three times as many large telescopes in the Northern Hemisphere as in the Southern, and 20 times as many professional astronomers in the north as below the equator. This situation, which was even more lopsided 30 years ago, is cited in the *Monthly Notes of the Astronomical Society of Southern Africa* from an article by A. D. Thackeray in the May issue of the *South African Journal of Science*.


The Great Aurora of September 22-23

JAMES E. McDONALD, *Institute of Atmospheric Physics, University of Arizona, Tucson*

AN INTENSE and widespread auroral storm began early in the evening of Sunday, September 22, 1957, and lasted until the predawn hours of the following day. At the peak of its activity, near 10 p.m. Mountain standard time, auroral observations were being made from coast to coast and all of the way from the Mexican border up to the Canadian line and possibly still farther north. In this summary all times cited are MST.

The diffuse reddish glow, typical of relatively rare low-latitude aurorae, was seen in Tucson, Arizona, at about 7 p.m. and then disappeared, only to reappear near 10 p.m., the time of nationwide maximum intensity. R. S. Shaw, of Tucson, has provided an excellent report of visual observations from the northern edge of the city, where artificial lights were nearly absent. A coppery red glow, from northwest to almost due east, was visible at about 9:45 p.m. above the Santa Catalina Mountains, some 10 miles away. Suddenly, at about 9:50, within only a few seconds the upper edge of this glow very rapidly ascended to an altitude of over 35 degrees, as estimated from Polaris, and remained there until the display faded near 10 o'clock. Occasional yellowish-white vertical rays emanated from near magnetic north, but apparently did not march laterally as such rays frequently do.

During inquiries to determine whether this display was limited to our Arizona



In the author's map of the great auroral storm of September 22-23, 1957, crosses mark sightings at 10 p.m. Mountain standard time; circles are stations not reporting aurora. The north-south elongation of the zones of visibility may indicate the influence of the earth's magnetic field.

area, I ultimately learned of a source of information of which I had been previously unaware—the informal special reports of aurora observations, labeled by the codeword AURBO, which are encoded at the end of the hourly airways weather reports, disseminated by teletype throughout the country. My discovery of this source of data was unfortunately belated, and on the evening of the 23rd it took a

diligent search through two large waste barrels, with the energetic aid of J. C. Killian of the Civil Aeronautics Administration's Tucson office, to retrieve the key teletype sequences containing the auroral data.

After filtering out sought-for sequences from bits of lunches and reams of yellow paper, only to discover that I had failed to hunt for the 11 p.m. data, I hesitated to ask Mr. Killian's renewed help in once again dumping and combing through the barrels. My reluctance, however, was quickly removed by his accusation that I seemed not to be showing the scientist's proverbial perseverance in getting the data at all costs, and together we found the missing hour's records in the depths of the well-filled barrels. The complete set of reports has permitted a reasonably full examination of the time development and the geographic coverage of this particular storm, so the scavenging seems now to have been worth while.

The accompanying map shows the pattern of auroral observations relayed at 10 p.m. MST by Weather Bureau and CAA stations at a total of about 260 points over the country. The teletype data available in the Tucson office includes most stations in the West but only a fraction of the eastern ones. This unevenness happened to be not too serious in the present case, since a very large area of overcast skies lay behind a cold front extending from southern Texas, along the Appalachians, and into New England; only scattered stations in that area could see



At the Sacramento Peak Observatory in New Mexico, Guenther Schwartz photographed the September 22nd aurora, using a Baker super-Schmidt meteor camera, only a portion of its 55-degree field being shown here. This 15-second exposure on Tri-X film was made at 10:49 p.m. Mountain standard time. Until then, trees masked the aurora, low in the northeast. "The arc that was visible was very reddish in color, the rays primarily whitish with a green tinge," Mr. Schwartz noted in his record book. Air Force Cambridge Research Center photograph.

the aurora anyway. By contrast, the western states were dominated by the sky-clearing action of the subsidence of air in a large anticyclone centered over Colorado on the evening of the 22nd; thus the West enjoyed a good opportunity for viewing this large auroral storm. The first teletyped accounts of auroral sightings came at 8 p.m., an hour after the earliest observations in Tucson, reached a maximum of 73 stations reporting aurora at 10 p.m., fell a bit near midnight, recovered slightly at 1 a.m., and ended after 4 a.m. on September 23rd.

Crosses on the map show all stations where aurorae were reported, and circles indicate places for which I found reports of completely clear skies at 10 p.m., but where no notation of aurora was included. Unfortunately the regulations for reporting aurorae at airways stations are such that we cannot conclude that every circle on the chart marks a spot where aurora was positively absent at that time. As part of a further study of this storm, I am now

making special inquiries at some 20 selected circled stations. More information from them will help interpret the odd elongation of the zones reporting the 10 o'clock aurorae, an elongation that seems roughly to parallel the magnetic meridians.

Of the 73 stations recording aurorae at 10 p.m., the farthest south was Douglas, Arizona, in latitude about 31° north. Grand Island, Nebraska, holds the distinction of reporting aurorae at every hourly observation from 8 p.m. to the close at 4 a.m.

The daily routine by which this airways data is processed makes it extremely difficult to extract information such as here summarized, if one tries it more than about one day after an auroral storm. This is because these auroral reports are not entered on the easily processed punch cards which the Weather Bureau uses for standard data. Instead, they are entered by hand on a certain form (WBAN 10-B) that is ultimately filed in the archives by

station rather than by hour, thus precluding easy synoptic analysis. Happily, however, the data are in convenient form if secured from any airways station before the original teletype sheets are thrown away (a daily disposal problem at such stations).

To my knowledge, no systematic study of these hourly airways reports of aurorae is now being carried out. The IGY auroral program being conducted by D. S. Kimball and C. W. Gartlein with Weather Bureau co-operation is, for example, drawing on only some 85 principal Weather Bureau stations, whereas there are about 500 airways stations in the country. Despite the less carefully controlled conditions of observation and recording in the case of the airways reports, I do believe that these provide interesting opportunities for study by amateur astronomers who live where teletype stations are maintained. The current peak of solar activity makes this year and the next especially good for such efforts.

LETTERS

Sir:

Page 333 of the May, 1957, *Sky and Telescope* reported the work of John S. Rinehart on the ablation or wastage of meteoritic bodies during their flight through the earth's atmosphere. My studies of this problem are presented in the book *Foundations of Meteoritics*, Moscow, 1955, and in the German journal *Chemie der Erde*, 18, 56-88, 1956.

From the properties of individual specimens of the Sikhote-Alin and other meteoritic showers, the following rule was established: If a specimen shows the earmarks of stable orientation during flight (roughly conical shape, flow marks on the front which have about 1/10 the diameter of the meteorite), then the rear surface has one of two kinds of structure—either with fully developed but much larger flow marks, or having sharp-edged irregularities.

The significance of these two types comes from the fact that there is little or no ablation at the rear surface of a meteorite falling with stable orientation. Suppose that the original meteoritic body broke up during its flight, and that a fragment preserved constant orientation during its remaining fall. Then its rear surface, if part of the original exterior, would show the large flow marks of the original body. On the other hand, if the rear face of the fragment were a surface of fracture, it would show the sharp irregularities of the fracture.

This reasoning may be applied to the photographs of the Cabin Creek meteorite that were published with the *Sky and Telescope* account of Dr. Rinehart's work. The approximately conical shape and the flow marks on its front show that flight was stable. But the back bears typical flow marks that are much larger than 1/10

the specimen's size. Hence we conclude that the Cabin Creek meteorite is only a fragment of an original body two or three times larger, perhaps 90 inches in diameter, to judge from the size of the rear flow marks.

Also, this rear surface formed part of the exterior of the original body. Therefore this specimen gives an excellent opportunity for studying the change in chemical composition with depth below the original surface. Such analyses give information on the influence of cosmic radiation on the formation of isotopes, including helium of mass three, in the meteorite.

E. L. KRINOV

Committee on Meteorites
U. S. S. R. Academy of Sciences
Moscow, U. S. S. R.

Sir:

While searching for Comet Mrkos on August 4th, we observed a phenomenon apparently related to the green flash. The western horizon was cloudless but hazy, and the setting sun deep red. We turned a mounted 3-inch 21x terrestrial refractor on the sun as it neared the horizon. Owing to layering of the air, the sun's limb appeared slightly notched; as the sun set, pairs of corresponding notches moved upward across the disk, pinching off small "bubbles" from the top.

Each bubble gradually turned orange, yellow, and finally pale green before it faded out, the whole process taking up to three seconds. By comparison with the telescopic disks of Venus and Jupiter, we estimated that the bubbles were about half a minute of arc in diameter. The upper limb of the sun also showed a yellow-green fringe, considerably narrower than the bubbles. Neither bubbles nor fringe could be seen in hand-held 7-power binoculars.

The green color seemed real; the ob-

servations of Venus and Jupiter showed it was not due to chromatic aberration in the telescope, and the gradual nature of the color and brightness changes rules out an explanation by eye fatigue.

On the following night the sky was clearer at sunset, the sun was brighter, and nothing unusual was observed.

W. E. HOWARD, III,

and A. T. YOUNG

Harvard College Observatory
Cambridge 38, Mass.

Sir:

On page 494 of the August issue, Edgar Everhart describes sidereal drives for telescopes, and proposes the accurate gear train $1 \text{ r.p.m.} \times 20/66 \times 14/56 \times 34/37 \times 1/100$. The identical ratio can be obtained from the train $1 \text{ r.p.m.} \times 5/55 \times 17/37 \times 1/60$, which is easier to construct because it requires fewer gears. Both trains are available in standard Boston gears. My train has a 5-tooth pinion and 20-pitch change gears.

ROBERT M. AYERS

1374 S. Branch Pkwy.
Springfield, Mass.

Sir:

The caption under the photograph of Jupiter on page 397 of the June issue contains an error. The prominent feature at the upper left is a disturbance in the south tropical zone, and not the red spot hollow. Mr. Botham's photograph on March 8, 1956, was taken when the longitude of the central meridian was 224° (System II). On that date the following end of the dusky disturbance was at longitude 215°, while the center of the red spot hollow was at 301°. Hence the latter feature was too near the following limb to be seen on the photograph.

ELMER J. REESE

R.D. 2, Box 396
Uniontown, Pa.

Amateur Astronomers

KANSAS CITY CONVENTION MARKS LEAGUE'S FIRST DECADE

ROLL CALL at the opening session of the general convention of the Astronomical League, held at the University of Kansas City, on August 31-September 2, revealed a representation of 42 per cent (more than 50 societies) of the league membership. The 269 delegates who registered came from all parts of the country, from places as widely separated as New England and California, Florida and Oregon, Texas and Wisconsin.

The convention marked the 10th year of the league as a formal organization, although it had its roots in the first convention of amateur astronomers at the New York World's Fair in 1939. In her address as retiring president, Grace C. Scholz recalled a map of the United States displayed at the Philadelphia convention in 1947, when the league was first permanently organized. The map had 72 pins marking all known astronomy clubs at that time. The number of societies in the country is now over 200.

In the past decade the Astronomical League has grown in membership from 30 local organizations to better than 130. Starting with four regions across the northern part of the country, the league now has nine, and each year the general convention is held in a different region.

Miss Scholz announced that the 1958 gathering of the league will be held over the 4th of July weekend, at Cornell University, Ithaca, New York, in the Northeast Region. Although the site for the 1959 convention has not been finally selected, an invitation has been received from the Mountain Astronomical Research Society Region to meet in Denver for a joint conclave with the Western Amateur Astronomers.

The league's two regular publications continue to grow in popularity. The monthly bulletin is edited by Mrs. Jane Gann, of the Columbus Astronomical Society, and will henceforth be called *Reflector*, this being the winning name in the recent contest to rename the bulletin. The *Junior Astronomer* is published in Washington, D. C., by Benjamin Adelman. At Kansas City, a special arrangement was announced whereby, with the support of Norman Edmund, *Reflector* will be sent directly to all individual members of league societies. This is an important milestone in the growth of what Miss Scholz called "our most important internal activity."

In 1952, the league became affiliated with the American Association for the Advancement of Science, and on several occasions has held sessions and displayed exhibit material at AAAS meetings.

Other important activities include the league's book service, at present con-

ducted by Margaret Kobs, of Portland, Oregon, and the support of two affiliated organizations, the American Association of Variable Star Observers and the Association of Lunar and Planetary Observers. In 1951 the first presentation of the Astronomical League award was made to Albert Ingalls, of telescope making fame, and since then a number of distinguished amateur and professional astronomers have received the award; none was made this year.

Miss Scholz told the meeting that the MOONWATCH satellite-observing program provides an unprecedented opportunity for the amateur. About half of the more than 90 teams now set up in the United States have been established in league societies.

She proposed that the league secure funds for developing a library of tape and slide lectures, which would be available to member societies on loan. A first attempt at this program indicates that considerable editing of the tape is required, and that there must be careful co-ordination between the speaker's words and the lantern slides that would be a part of each lecture. The league officers are also seeking the co-operation of the National Science Foundation and the American Astronomical Society in procuring astronomers for lectures at local meetings and for workshops at colleges.

Concluding her address, Miss Scholz pointed out that, although funds for expansion are always important, the present greatest need of the league is for manpower—for leaders who are willing to accept responsibility and who will carry through on projects of benefit to local societies and amateur astronomy in general.

Russell C. Maag, of the Central Missouri Amateur Astronomers, was elected president of the league to succeed Miss Scholz, his term of office beginning at Kansas City.

The exhibits at general conventions continue to grow in importance, and the exhibit area forms a place where discussions between sessions become detailed and lengthy. The Kansas City society prepared a special display of international astronomy from many parts of the world, the material being gathered from societies in Europe, South America, South Africa, and Asia.

There were elaborate displays from several of the manufacturers who are supporting members of the Astronomical League; in several cases their chief officers personally attended the convention to demonstrate their instruments and to learn the needs of amateur observers. The Kansas City program listed 10 supporting members: Coast Instrument, Inc., Criterion Manufacturing Co., E and W Optical Co., Edmund Scientific Co., J. W. Fecker, Inc., S. E. Laszlo, Sky Publishing Corp., Skyscope Co., Spitz Laboratories, Inc., and United Scientific Co.

On Saturday afternoon, there was a satellite-observing program, with five invited speakers, including Col. Owen F. Clarke, U. S. Air Force liaison officer for MOONWATCH, who described the problems of flying airplanes to simulate satellites. Later that day, the delegates toured the Linda Hall science library, under the guidance of its director, Dr. Joseph Shipman. The library regularly subscribes to 7,000 periodicals in science and industry. That evening a star party and observing technique session was scheduled, but clouds covered the moon just 20 minutes before the predicted occultation of Saturn, so that event could not be observed by the delegates.

Sunday's schedule included a junior session and a talk on "Mathematics and As-



Members and guests at the general convention of the Astronomical

League



President Grace C. Scholz opening the Kansas City convention of the Astronomical League. Left to right are Gene L. Tandy, general chairman, Dr. Richard M. Drake, chancellor of the University of Kansas City, Mrs. Wilma Cherup, executive secretary, Miss Scholz, Dr. William C. Doyle, S. J., Rockhurst College, and Fred Hartman, president of the Astronomy Club of Kansas City.

tronomy" by Dr. William C. Doyle, S. J., of nearby Rockhurst College. The convention banquet was held there Sunday evening, with the principal speaker Dr. Richard N. Thomas, of the National Bureau of Standards, Boulder, Colorado. After describing briefly the current problems of the sun's chromosphere, Dr. Thomas appealed to the young people in his audience to take up with him their problems of how to follow astronomy as a career. As a result, later that evening he held a long informal discussion with the many junior astronomers attending the convention.

The Mid-States regional meeting was held Monday morning, and then a panel of experts, including Dr. Thomas, answered numerous written questions from delegates. Following the final convention

session, the 10th anniversary meeting of the Association of Lunar and Planetary Observers was called to order at 11 a.m. The society's director, Walter Haas, was chairman for a lively program of papers concerning the moon, Comet Arend-Roland, Mercury, Venus, Jupiter, and Mars. The afternoon program of the ALPO brought the Kansas City convention to a close.

C. A. F.

EATONTOWN, NEW JERSEY

The recently organized Society of Telescopy, Astronomy, and Radio (STAR) now has 33 members. Officers are Frank K. Priebe, president; Herbert D. Tanzman, treasurer; and Mrs. Thomas B. Richey, secretary, whose address is 2038 Magill Drive, Eatontown, N. J.

THIS MONTH'S MEETINGS

Cambridge, Mass.: Amateur Telescope Makers of Boston, 8 p.m., Harvard Observatory. Nov. 14, Dr. Gerald S. Hawkins, Harvard Observatory, "Comets."

Detroit, Mich.: Detroit Astronomical Society, 2:30 p.m., State Hall, Wayne University. Nov. 10, Dr. Helen W. Dodson, University of Michigan Observatory, "The Solar Program of the International Geophysical Year."

Madison, Wisc.: Madison Astronomical Society, 8 p.m., Washburn Observatory. Nov. 13, Dr. V. E. Suomi, "The IGY and the Space Satellite."

New York, N. Y.: Amateur Astronomers Association, 8 p.m., American Museum of Natural History. Nov. 6, Dr. Ernest G. Reuning, University of Pennsylvania, "An Astronomer Looks at Space Travel."

Washington, D. C.: National Capital Astronomers, 8:15 p.m., Commerce Department auditorium. Nov. 2, Dr. S. Fred Singer, University of Maryland, "Meteors."

NEW YORK JUNIORS

Now entering its 29th season, the Junior Astronomy Club of New York is offering five astronomy classes, in addition to its regular lecture series. The courses cover elementary and intermediate astronomy, and also mathematics and electronics as applied to astronomy.

Students in the electronics course hope to construct such instruments as a photometer, an interferometer, and a photoelectric clock drive devised by Stephen E. Strom, club president. The clock drive would operate a telescope automatically.

Membership information may be obtained from Mr. Strom, 1950 Andrews Ave., Bronx 53, N. Y.

PATERSON, NEW JERSEY

The North Jersey Astronomical Society, now in its third year, has a membership of eight adults and 10 juniors. Most of the members own their own telescopes, and the club has made a 6-inch reflector.

In addition to holding public star parties at Paterson State Teachers College and other sites, the society is currently setting up a MOONWATCH station at East-side high school, under the supervision of R. DelVecchio. Visual aurora observations during the International Geophysical Year are being undertaken.

Officers are Wilbert M. Diggles, president; Eleonore Kraus, treasurer; and Donald Trombino, secretary. Further information may be had from Mr. Trombino, 4-4D Alabama Ave., Paterson 3, N. J.

SANTA ANA, CALIFORNIA

The permanent meeting place for the Orange County Amateur Astronomers is now room U12 of Santa Ana College. The club meets monthly at 7:30 p.m. on the first Friday, according to the secretary, Glenn H. Shaw, 2061 Halladay, Santa Ana, Calif.



League, University of Kansas City, August 31-September 2, 1957.

The Most Massive Stars Known

OTTO STRUVE, *Leuschner Observatory, University of California*

A VITAL PART of modern astrophysics had its beginning about 150 years ago, when it was realized that many visual double stars were not mere chance pairs but actual binary systems, often with conspicuous orbital motion. This motion was the visible effect of the gravitational attraction of one star upon the other. Since that time, astronomers have succeeded in determining the masses of the component stars of many binaries.

Data on star masses is fundamental to our newly gained understanding of stellar structure and stellar evolution, yet, even today, the only sure basis of this data is the orbital motion of the visual and spectroscopic binaries. We cannot measure the mass of an isolated star—except

the sun, which has planetary companions. We are forced to assume, as a first approximation, that the masses of single stars resemble those of double star components having similar luminosities, diameters, and surface temperatures.

Popular writers usually avoid explaining how the masses of double stars are computed because the problem involves a fair amount of algebra if all kinds of binaries are included in the discussion. But the basic ideas are quite simple, and the masses of the heaviest known stars can be found from elementary considerations.

Large stellar masses are of particular interest. What is the heaviest star or pair of stars now known? Is there a physical limit to the amount of matter a star can accu-

mulate? And what are the properties of the heaviest stars?

We choose for our example Plaskett's star, a very massive binary system just visible to the naked eye, located in the constellation of Monoceros about midway between Procyon and Betelgeuse. Also known as HD 47129, this object is of visual magnitude 6.1, and its spectrum is of type O8. In 1922, the Canadian astronomer J. S. Plaskett discovered it to be a spectroscopic binary, from spectrograms taken with the 73-inch reflector of the Dominion Astrophysical Observatory at Victoria, British Columbia. The period of orbital motion is 14.4 days.


In Fig. 1 are plotted observations of the radial velocities of the two stars that constitute this system. The velocity curve of the primary star—the one with the stronger absorption lines—is represented by the dots. The symmetry of the curve shows that the orbit of this star relative to the other is very nearly circular.

What we measure on the spectrograms are the Doppler displacements of the absorption lines at different times (see Fig. 3). This gives us the star's radial velocity, that is, the component of the true velocity that lies along the line of sight, at each phase. At phase 0 day, the primary star is approaching us with a velocity of 190 kilometers per second, and at phase 7.2 days it is receding at 240 kilometers per second.

A part of the velocity is due to the motion of the system as a whole relative to the sun. Were this motion zero, the circular orbit of the primary star around the center of gravity would produce equal values for the maximum observed velocities of recession and approach. The fact that these two velocities are unequal shows that this is not the case—the center of gravity is receding from us with a constant velocity of $\frac{1}{2}(240 - 190) = 25$ kilometers per second. The dashed line in Fig. 1 indicates this motion of the center of gravity.

Thus, with respect to the center of gravity, the primary star is moving $190 + 25$, or 215 kilometers per second when its velocity of approach is greatest, and $240 - 25$, or 215 kilometers per second at maximum velocity of recession. This would be the orbital velocity of the primary star with respect to the center of gravity, provided the orbital plane were in the line of sight, that is, if the orbit were presented to us edgewise.

But since HD 47129 is not an eclipsing



Spectroscopic observations of stars, including binary systems, form the principal work of the 73-inch reflector. The instrument was built in 1918, and is housed here, at the Dominion Astrophysical Observatory, near Victoria on Vancouver Island, British Columbia.

variable star, we infer that the orbit is not edge-on; there appear to be no rhythmic light variations caused by one star eclipsing the other at each revolution. The true orbital velocity must be larger than 215 kilometers per second, being diminished by foreshortening. Exactly how much larger is not known, but we shall probably not be far from the truth in assuming that the real, unforeshortened velocity is about 250 kilometers per second. This value corresponds to a tilt of the orbit plane of 30 degrees to the line of sight.

Assuming, then, that the primary star of HD 47129 is moving in a circular orbit at 250 kilometers per second, what can we deduce concerning the masses in this system? We shall begin by applying Newton's first and second laws of motion. The first law is that a moving body not subjected to any external force will continue to move in the same direction and with unchanged velocity. Newton's second law states that a body moving with changing velocity or direction is being subjected to a force that is equal to the product of the body's mass and its acceleration (the change in amount and direction of velocity). This law is expressed in the well-known formula, $f = ma$.

Suppose, in Fig. 2, that the primary star is in position *A*. Its orbital velocity of 250 kilometers per second is tangential to the circular orbit, as shown by the arrow. If no force were acting on the star, it would arrive at *D*, the end of the arrow, after a lapse of one second. In reality the star arrives at position *B*. Its speed is still 250 kilometers per second, but the direction of the next velocity arrow, starting from *B*, is not the same as that of the initial velocity at *A*. The star has experienced a central force that has caused it to "fall," in the course of one second, from *D* to *B*.

This distance of fall in one second is numerically just half of the acceleration. It is like the case of a stone falling from rest to the surface of the earth. Its acceleration or increase in velocity is 32 feet per second each second; and it falls 16



Plaskett's star is named after the Canadian astronomer J. S. Plaskett, director of the Dominion Astrophysical Observatory from 1917 to 1935.

feet in the first second. If the stone has an initial horizontal velocity, the result is still the same, its fall toward the earth in the first second is still 16 feet, or half the acceleration.

In Fig. 2, one leg of the triangle *CAD* is the radius of the orbit, the distance between the center of gravity and the star. The circumference of the orbit is the orbital velocity (250 kilometers per second) times the period (1.24×10^6 seconds), or 3.1×10^8 kilometers; the radius is thus about 5×10^7 kilometers. The other leg of the triangle, *AD*, is the star's motion in one second, or 250 kilometers. Finally, the hypotenuse is the orbital radius, 5×10^7 kilometers, plus the distance of the fall, which we shall call *x*.

From the Pythagorean theorem for a right-angle triangle,

$$(5 \times 10^7 + x)^2 = (250)^2 + (5 \times 10^7)^2.$$

Because x^2 is relatively very small we can neglect it, and a little arithmetic shows

that $x = 6.25 \times 10^{-4}$ kilometers, or 62.5 centimeters. Doubling this distance of fall gives for the acceleration 125 centimeters per second each second. Newton's second law can therefore be written $f = 125m$, where *f* and *m* are in dynes and grams, respectively, the units in the centimeter-gram-second system.

In order to solve for the mass, however, we need Newton's law of universal gravitation. This states that the force in dynes between any two bodies is equal to 7×10^{-8} (the gravitational constant) times the product of their masses in grams, divided by the square of their distance apart in centimeters.

But we know only the primary star's distance from the center of gravity, which can be called *a*; for the other star the corresponding distance is *a'*, and the stars' total separation is $a + a'$. From Fig. 1 we may infer, despite the large scatter in the measurements, that the secondary's velocity curve has about the same range as that of the primary star. Therefore the radius of the secondary orbit should be about the same as the primary's, and *a'* should also be about 5×10^7 kilometers. The sum of these distances, $a + a'$, is thus twice this value, or 10^{13} centimeters.


Now we are ready to equate the force, which acts on the primary star according to the law of gravitation, with the force expressed by the second law of motion, making the assumption that gravitation alone holds the two stars in their orbits:

$$(7 \times 10^{-8})mm'/(a + a')^2 = 125m.$$

The factor *m* can be cancelled from both sides of this equation, while $(a + a')^2$ is 10^{26} . The equation can be solved for *m'*, the mass of the secondary star, which comes out 1.8×10^{35} grams.


Since the mass of the sun is 2×10^{33} grams, the secondary component of Plaskett's binary is about 90 times as great. The primary star should also be about 90 solar masses, because we believe that the center of gravity is about midway between the stars. The total mass of this double star is thus some 180 suns!

It is important to review our assumptions. We supposed that the force deduced from the observed acceleration is purely gravitational, but in principle it is



Left: Fig. 1. K. D. Abhyankar, of Leuschner Observatory, has plotted radial velocity observations of HD 47129, using dots for the primary star, circles for the secondary, from spectrograms taken at Victoria, McDonald, and Lick Observatories.

Right: Fig. 2. In this diagram, where the circle represents the orbit of the primary star about the center of gravity, *x* indicates the distance of fall in one second.






Fig. 3. In spectra of HD 47129, the diffuse dark lines of the primary star are shifted by the Doppler effect. Sharp dark lines are of interstellar origin, hence undisplaced by orbital motion.

the sum of physical forces of all possible types. Yet we know that in the solar system gravitation by itself suffices to explain the motions of the planets and satellites — magnetic, electrical, and other forces are negligible. If they were of consequence in a double star system, they would be detectable from their spectroscopic effects.

Another assumption involved the tilt of the orbital plane to the line of sight. It is pretty certain that we have not underestimated the true orbital velocity of the primary component. As has been mentioned, if the orbital plane were close to the line of sight, Plaskett's star would be an eclipsing binary, but no eclipses have been observed. If the orbit plane is tilted to the line of sight by more than the 30 degrees we selected, then the true orbital velocity would be even greater than 250 kilometers per second, and the resulting masses would be larger. Therefore, this assumption yields minimum values for the masses.

Our third assumption was that the two components have equal masses. This rests upon the rough estimate from Fig. 1 that the range in radial velocity is the same for both components. If, as seems likely, the range of the secondary is somewhat smaller than the primary's, then the secondary is the more massive star.

These stellar masses, of the order of 90 times the sun's, are the largest definitely known at the present time. We may therefore conjecture that the upper limit of possible star masses is about 100 times the sun, or 2×10^{35} grams. It was suggested many years ago by Sir Arthur Eddington that the radiation pressure inside a star of this mass is high enough to offset the internal gravitation that holds the star together — if there were any star of larger mass, radiation pressure would blow it apart.

It is therefore of interest that Plaskett's star appears highly unstable. One or both components are shedding gas into interstellar space, and in many respects behave like a slowly exploding star of the P Cygni or Wolf-Rayet type.

Fig. 3 shows a portion of the spectrum of Plaskett's star at two phases, 7.9 and 12.3 days. The large Doppler shift of the primary component can easily be seen for the helium line at 4026 angstroms. Features arising from the secondary star can hardly be discerned in this spectral region, the ultraviolet, but are easily seen in the red. The very sharp dark lines marked K and H are of interstellar origin; they result from the absorption of light by

ionized calcium atoms situated between us and the star.

Of special importance is the strong line of helium at 3888 angstroms, discovered by J. Sahade. This absorption line is present only at phases close to 7.9 days, and then gives a velocity of approach of about 700 kilometers per second. At that time there is something resembling a rapidly approaching jet of helium gas which passes between us and the primary star. It is believed that this jet originates from the secondary component, and diffuses toward all sides to form ultimately a vast expanding shell of tenuous nebulous

matter. The primary component also ejects gas, especially hydrogen, but fairly uniformly in all directions.

Other evidence of the instability of Plaskett's star is provided by some of the absorption and emission lines, which suffer irregular changes from cycle to cycle. It is almost certain that similar variations will be found in the brightness of this system. Because of its instability, the star should be observed photoelectrically. In many respects, Plaskett's star is among the most remarkable known, and it promises us a rewarding insight into the problems of very massive stars.

Double Star Studies in Indonesia

At the Bosscha Observatory, near Bandung, Java, photographic observations of double stars form the main program of the 23½-inch refractor. At this nearly equatorial station in Indonesia, it is possible to reach practically all of both the northern and southern skies under very favorable atmospheric conditions.

In his report for the years 1955 and 1956, Director G. B. van Albada tells of special studies of several well-known double stars, including Sirius, Antares, Albireo, and Alpha Centauri. He also reports on the photometry of galactic clusters and observations of Mars in three colors during its 1956 opposition.


In a recent paper concerning photographic observation of binary systems with very large magnitude differences, Dr. van Albada describes the special techniques he has developed to obtain measurable photographs of such difficult pairs as Sirius, with results like that pictured here. The companion is seen above and to the left of the overexposed central image of Sirius. From the astrometric viewpoint, the faint diffraction images on either side of the central one are especially important.

It is standard procedure in double star photography, if the primary star is several magnitudes brighter than the secondary, to produce such fainter images by means of a coarse wire diffraction grating placed in front of the telescope objective. These images are actually spectra of the first order, second order, and so forth. Because one pair of them are of about the same size and density as the image of the companion, the latter's position with reference to them may be measured much more accurately than with respect to the overexposed central image of the primary.

Difficulties arise with this method, however, if the companion is very much

fainter than its primary, as in the case of Sirius. Dr. van Albada has found that a grating of only five or six parallel wires, widely but *unequally* spaced, produces the desired result. The spiked appearance of the central image is caused by a hexagonal diaphragm in front of the objective, which produces a strong diffraction of light but reduces the intensity of the image in the spaces between the spikes. The diaphragm is oriented so the image of the companion falls in one of the clear spaces.

The Bosscha Observatory report discusses Albireo (Beta Cygni) as a triple system. Visually, this famous double star has a separation of about 35 seconds of arc. But measurements of photographs in blue and yellow light give positions differing by 0.14 second for the brighter component, which is known to have a composite spectrum. Analysis of all available observations suggests strongly that the yellow component of the unresolved pair is moving in an elliptical orbit around its blue neighbor.



In this recent photograph of Sirius taken with the 23½-inch refractor, the famous white dwarf companion appears above and to the left of the overexposed image of the primary star, which is flanked by diffraction images that can be measured for position. The enlargement from the original negative is about 13 times. From Bosscha Observatory "Contributions" No. 3.

OBSERVER'S PAGE

Universal time is used unless otherwise noted.

THE LUNAR ECLIPSE OF NOVEMBER 7TH

THE last of the four eclipses of 1957 will be a total eclipse of the moon on the morning of Thursday, November 7th, whose opening stages will be visible in the western parts of the United States and Canada. For amateurs living east of the Mississippi River, generally speaking, the moon will have set before the eclipse commences. At most places along the Pacific Coast, however, our satellite will be totally eclipsed by the time of moonset.

The timetable that follows has been adapted from the *American Ephemeris and Nautical Almanac*; all times are a.m.

	CST	MST	PST
Moon enters penumbra	5:30	4:30	3:30
Moon enters umbra	6:43	5:43	4:43
Total eclipse begins	8:12	7:12	6:12
Middle of eclipse	8:27	7:27	6:27
Total eclipse ends	8:42	7:42	6:42
Moon leaves umbra	10:10	9:10	8:10
Moon leaves penumbra	11:23	10:23	9:23

As seen from St. Louis, Missouri, the moon will set 10 minutes before umbral eclipse begins; at Minneapolis, Minnesota, moonset comes 16 minutes after the umbral shadow starts to take its first "bite" out of the moon. Only very limited observations can be planned for such points. But farther west, more and more of the eclipse may be seen. At Denver, Colorado, the moon sets 35 minutes before the commencement of totality, and at Tucson, Arizona, 24 minutes before.

West Coast watchers will be the most favored of any in the United States, particularly in more northerly latitudes, where moonset will be delayed because of the north declination (+16°) of the moon. At Pasadena, California, eight minutes of total eclipse may be observed before the moon descends below the western horizon. For Berkeley, California, moonset comes at 6:45 a.m., Pacific standard time, three minutes after the end of totality. At Port-

land, Oregon, the geometrical conditions are even better, as 20 minutes may be seen of the partial stage following totality.


For early-rising amateurs who wish to make visual or photographic observations of the phenomenon, there are many practical suggestions in previous issues of *Sky and Telescope*. Consult page 43 of the November, 1956, number for detailed information on emulsions, lens speeds, and exposure times for filming the eclipse. Page 38 of that same issue lists some projects for visual observers. Among the suggestions made are descriptions of colors on the eclipsed moon, notes on the visibility of craters and seas inside the umbral shadow, and careful records of the times when individual craters enter or leave the umbra.

Even with the unaided eye, a lunar eclipse is an interesting phenomenon to watch. In some places, the eclipsed moon and the sun may be seen above the horizon simultaneously, a seeming paradox explained by atmospheric refraction, which apparently raises both bodies. Binoculars are worth using, and refractors of 2- to 4-inch aperture are well suited for visual observing, as are small reflectors.

The next lunar eclipse will be the partial one of May 3, 1958; like this year's event, it will be visible for United States observers only from the western states.


"THE PHOTOELECTRIC PHOTOMETER"

A wide, inviting field of observational work is open to the advanced amateur who builds a photoelectric photometer and uses it on his telescope for brightness measurements of variable stars. The basic requirements are some practical acquaintance with electronics, a well-mounted telescope of moderate size with a good drive, and a certain amount of astronomical experience. For several years the American



Events on this chart of the moon's path are labeled in Universal time.

Follow the Stars Wherever You Go



with a BALscope TELESCOPE

Now you can enjoy astronomy whenever you wish—wherever you are, thanks to the excellent portability of the BALscope 60mm telescope. Why, it's so compact (16½" long, weighs 48 ounces) you can carry it anywhere, and it can be mounted on any pan-head camera tripod with a B&L tripod adapter.

Focusing the BALscope is simple yet precise with its unique prism-focusing system. Its high light-gathering power and lack of stray light enables you to observe stars down to the tenth magnitude on moonless nights. Four easily interchangeable eyepieces ranging from 15X to 60X provide excellent results—from Star Clouds to Double Stars. Of course, you'll find the BALscope is also the finest terrestrial telescope available.



You can own the BALscope for just pennies a day under new Time Payment Plan offered by B&L dealers everywhere.

\$6.18 per mo.
\$115 cash price

WRITE for free copy of G-36, "The Telescope." Tells all about the BALscope and how it is so ideally suited to amateur astronomy. Tells also of many ways your family will enjoy the BALscope. Bausch & Lomb Optical Co., 25823 Lomb Pk., Rochester 2, New York.

BAUSCH & LOMB



THE MARK III Sidereal Telescope Drive

Bodine Synchronous Motor, Automatic Clutch, Automatic Sidereal Time Circle requiring no calculation for location of objects, R.A. Indicator, Sidereal Time Indicator, Lifetime Quality, Precision Fabrication.

Ramsden Eyepieces
Declination Circles
Hour-Angle Circles

Write for price list

H & W OPTICAL CO.
654 Milwood Ave., Venice, Calif.

Astronomy Films

16-mm. sound, 400-foot reels
I THE SUN; II THE MOON;
III LAR SYSTEM; IV MILKY WAY;
V EXTERIOR GALAXIES.

2 x 2 SLIDES

35-mm. STRIPS OF SLIDES

THROUGH 200-INCH AND
OTHER GREAT TELESCOPES

Catalogues on request.

International
Screen Organization

1445 18th Ave. North, St. Petersburg 4, Fla.

TRIPLES 'SCOPE PERFORMANCE!

Sharper images, wider field, more light at higher powers! A startling statement positively proven in 16-page telescopic educational matter, plus many helpful hints, sent free on receipt of self-addressed long envelope bearing 9c return postage.

First, the Goodwin Resolving Power lens placed in front of eyepiece gives three times the magnification on each by narrowing your primary focal length angle up to three times, yet extends eyepiece out no more than two inches from normal. This alone sharpens definition.

Next, by achieving your highest powers on more comfortable low-power eyepieces, you lessen image deteriorations due to short-focus acute bending of the convergent beam, since all usual eyepieces are $f/1$ or less. Again sharper images from this highest precision lens.

Third, you get greater illumination and wider field by relieving tiny aperture restrictions of higher-power eyepieces.

The Resolving Power lens is achromatic, coated, gives flat field sharp to the edge. Here is astonishment! Price \$23.50 in 4" long adapter tube fitting standard $1\frac{1}{4}$ " eyepiece holders ONLY. (Also adaptable to Unitrons; state if Unitron.) Money back if not positively thrilled after two weeks trial! Used and praised by legions!

No COD's—Colleges and Observatories may send purchase order.

FRANK GOODWIN
345 Belden Ave., Chicago 14, Ill.

Association of Variable Star Observers has had a committee under the chairmanship of Lewis J. Boss to encourage photoelectric work.

This committee has published a small mimeographed manual, *The Photoelectric Photometer*, which contains instructions and circuit diagrams for the construction of a simple photometer and amplifier. There are brief directions for its operation and for the reduction of measurements made with it, based on the successful amateur work of J. J. Ruiz, Danemora, New York. This manual is available at \$2.00 per copy from Mr. Boss, 2111 28th Ave., San Francisco 16, Calif.

SEEING PLANETS BY DAY

Several years ago I planned to observe all of the bright planets by day. Mercury, Venus, Mars, and Jupiter were so seen, but I was not successful with Saturn until August 16th of this year. Using 6 x 30 binoculars, I spotted this planet in a clear sky six minutes before sunset. Since the predicted magnitude of Saturn on this date was +0.7, this means that bright stars like Sirius and Canopus should not be too difficult to see in full daylight.

JOSEF R. OTOUPALIK
800 $\frac{1}{2}$ Ninth St.
Greeley, Colo.

SUNSPOT NUMBERS

These are observed mean relative sunspot numbers from Zurich Observatory and its stations in Locarno and Arosa.


August 1, 150; 2, 148; 3, 178; 4, 166; 5, 147; 6, 162; 7, 167; 8, 141; 9, 121; 10, 88; 11, 95; 12, 118; 13, 120; 14, 135; 15, 170; 16, 198; 17, 189; 18, 197; 19, 185; 20, 170; 21, 144; 22, 147; 23, 114; 24, 104; 25, 138; 26, 164; 27, 182; 28, 222; 29, 244; 30, 255; 31, 282. Mean for August: 162.6.

September 1, 257; 2, 230; 3, 201; 4, 166; 5, 184; 6, 160; 7, 137; 8, 175; 9, 250; 10, 265; 11, 255; 12, 264; 13, 260; 14, 263; 15, 265; 16, 283; 17, 258; 18, 295; 19, 317; 20, 294; 21, 334; 22, 302; 23, 268; 24, 239; 25, 234; 26, 220; 27, 227; 28, 249; 29, 249; 30, 229. Mean for September, 244.3.

The following American sunspot numbers for July and August were derived by Dr. Sarah J. Hill from AAVSO Solar Division observations:

July 1, 181; 2, 166; 3, 198; 4, 199; 5, 206; 6, 197; 7, 153; 8, 144; 9, 129; 10, 119; 11, 98; 12, 76; 13, 106; 14, 113; 15, 141; 16, 148; 17, 175; 18, 186; 19, 189; 20, 211; 21, 220; 22, 252; 23, 227; 24, 189; 25, 171; 26, 166; 27, 155; 28, 138; 29, 110; 30, 112; 31, 97. Mean for July, 160.4.

August 1, 102; 2, 126; 3, 132; 4, 126; 5, 116; 6, 142; 7, 125; 8, 126; 9, 114; 10, 90; 11, 97; 12, 95; 13, 109; 14, 133; 15, 163; 16, 173; 17, 175; 18, 175; 19, 173; 20, 125; 21, 112; 22, 95; 23, 86; 24, 99; 25, 145; 26, 144; 27, 167; 28, 191; 29, 210; 30, 236; 31, 228. Mean for August, 139.7.




THE LUNAR CRATER PLATO

Jackson T. Carle's article "Three Riddles of Plato," in the April, 1955, *Sky and Telescope*, encouraged me to begin observations of the details on the floor of this crater, using an 8-inch $f/7.5$ reflector.

Sketches on several nights of poor seeing showed only a few of the interior craterlets. But on August 12, 1957, seeing was superior when the accompanying drawing was made at 11:30 p.m. Central standard time. The terminator of the waning moon passed across Mare Crisium then, the sun's colongitude being 120°7.

No fewer than 26 craterlets and small bright spots inside the crater were drawn. Most of them were found with 200x, and others were detected with 400x and 600x. For one brief instant of perfect seeing, the dark velvety floor of Plato appeared to be covered with still other minute spots.

PATRICK McINTOSH
R.R. 2, Box 231, Robinson, Ill.




THE ASTRONOMICAL LOCATOR

is an instrument of many uses to teach or to study the solar system.

- By turning the large 12-inch disk with its apparent sun globe to any hour on the small dial, one can see readily what part of the sky is currently overhead and which constellations along the ecliptic are in view at night.
- The equation of time, with the months in which it must be used, is near the earth globe to show in minutes what adjustment to the sun or to the clock is necessary.
- Learn when and where to look for specific stars with the aid of this instrument. \$36.00 postpaid

PREMIER PLASTICS
OF MILWAUKEE

204 W. WASHINGTON ST., MILWAUKEE 4, WIS.



QUESTAR GUARANTEES 0.9-SEC. RESOLUTION ON NATIONAL BUREAU OF STANDARDS TEST CHART. FINAL TEST RECORD AND CHART COMES WITH EVERY INSTRUMENT.

If telescopes are guaranteed at all, as they ought to be, they are usually guaranteed to equal their theoretical resolving power. For Questar's $3\frac{1}{2}$ -inch aperture the Dawes limit is 1.3 seconds of arc. Since nature seems to favor small apertures of some 3 to 5 inches, a telescope in this size range is poor indeed if it cannot meet its theoretical limit of resolution. It therefore seems to us that such a guarantee means little. Further, it puts the burden of proof entirely on the purchaser.

Absolutely tremorless and quiet air is essential to the testing of a telescope if you really want to know how well it will perform. When an instrument is tested on the night sky, the rays of light that reach it must traverse the earth's entire atmosphere. The least movement in the disc and ring system of a star image is evidence of atmospheric interference, as W. H. Pickering has pointed out. Bell tells of a famous English astronomer who "had seen but one first-class night in 15 years." One Questar owner observed for two years before a night of exceptionally fine seeing revealed his instrument's full capabilities. This all sounds discouraging, but we are

speaking of not just good seeing, but of seeing that is perfect.

We can make our own perfect seeing by drastically shortening the air path through which we look to read the proper test chart, and this will demonstrate precisely the resolving power of a telescope in seconds of arc. The charts we use are purchased from the Bureau of Standards. They may be used outdoors in the quiet air of early morning or late afternoon at distances up to 300 feet, or even indoors at only 41 feet.

Here is a method of testing telescopes on which we can base a performance guarantee and at the same time furnish the simple means to prove it. We do not ask the novice to familiarize himself with star or planetary test objects, and then wait weeks or months for steady air to look through. The man who buys a Questar now may have room to test it indoors the very day it arrives, regardless of the wind or rain or cold of outside weather. He can duplicate our test himself; the chart will tell him at a glance whether his $3\frac{1}{2}$ -inch Questar resolves the 0.9 second of arc that we guarantee.

In going to the chart system, we only do what the Air Force and other photographic people have been doing for years. They have to know the exact quality of some very large and costly lens systems, and various kinds of charts and patterns for this purpose are in common use.

In consulting the experts at the Bureau of Standards on our special needs, they suggested that we use a chart which they make photographically on glossy paper, the smallest lines of which cannot be seen with the naked eye. The Bureau's recommendations for its proper use are embodied in the directions which we send with every chart.

The De Luxe Questar sells complete with all accessories at \$995. The Field Model at \$495 consists of the same major optics in a simple barrel without built-in auxiliary devices, with bracket casting for mounting on a panhead tripod, 40x eyepiece and choice of star diagonal or erecting prism. This model will accept $1\frac{1}{4}$ -inch standard eyepieces.

QUESTAR CORPORATION
New Hope, Pennsylvania



Excitingly New!

CRITERION'S STANDARD DYNASCOPES

6" - 8" - 10" - 12" - 16" starting as low as \$265

Custom built to meet a professional's exacting specifications

Priced within easy reach of the serious amateur

Only Criterion could produce such magnificent instruments at such reasonable cost. Combining the latest advances in optical engineering with the old-world skill and patience of master craftsmen, these superb new Dynascopes, custom made, offer top-quality performance. Every feature necessary for superior viewing is precision finished to professional standards. Optical surfaces accurate to 1/10 wave.

The five matched eyepieces are the finest orthoscopic and achromatic. Mechanical features are carefully custom built to assure trouble-free operation. Construction throughout is of lifetime materials. There is both maximum rigidity and easy portability.

Above all, Criterion's strict quality control allows no room for chance. Each Dynascope is doubly tested and inspected; these rigid tests must be flawlessly met before shipment. Your satisfaction UNCONDITIONALLY guaranteed. Yet the price is amazingly low.

6" Standard Model (80 lbs. shipping weight f.o.b. Hartford) **\$265**

Also illustrated: Electric Clock Drive, \$80; Setting Circles, \$60; Permanent Pedestal, \$70.

Write today for full specifications and details of these superior reflecting telescopes. Easy payment terms available on request.


Superior Features Include

- f/9 Parabolic Mirror, accurate to 1/10 wave
- Fully Rotating Tube, for comfortable viewing
- Massive Equatorial Mount, adjustable for your latitude
- 5 Matched 1 1/4" Eyepieces — 3 Achromatic Ramsdens (45X, 76X, 152X) — 2 Orthoscopes (228X, 343X)
- 8 x 50 Achromatic Finder Scope with cross-hairs
- Secondary Support that minimizes diffraction
- Declination Axle, 1 1/2" diameter, with slow motion
- Polar Axle with Needle Bearings on 1 1/2" shaft
- Rack-and-pinion Focusing
- Double-draw Focusing Tube for any eyepiece
- Precision-fashioned 54" Bakelite Tube
- Exclusive Two-way Pier-tripod — a massive 45-pound pier for permanent installation
- Lightweight Tripod hidden inside pier for field trips

CRITERION MANUFACTURING COMPANY

Manufacturers of Quality Optical Instruments

Dept. D-9, 331 Church St., Hartford 1, Connecticut



Positions of the eight deep-sky objects described below are plotted on this chart, which shows stars to apparent visual magnitude 7.75.

DEEP-SKY WONDERS

AQUARIUS, a large constellation well placed for observing this month, contains a variety of objects for the amateur with a not-so-small telescope. Their locations are shown on the accompanying chart prepared from the Skalnate Pleso *Atlas of the Heavens*, which is useful for this study, as it contains many objects not included on older charts.

NGC 6981 (M72) is on all the charts, for the Messier clusters and nebulae are among the brightest in the sky. This globular cluster, situated at right ascension 20^h 50^m.7 and declination -12° 44' (1950 co-ordinates), is five minutes of arc in diameter; its total light matches that of a 9th-magnitude star. In a 10-inch reflector it appears as a dense swarm of faint stars. Nearby on some maps is the non-existent cluster M73, actually only a grouping of four stars. Those who still write that they can discern a nebula or cluster here might well consult Admiral Smyth's *Celestial Cycle*, where, in his account of M72, he wrote over a century ago that M73 does not exist.

NGC 7009, the Saturn nebula, curiously is not on the Messier list, although it is one of the show objects of the sky. On an extra fine night look for it at 21^h 01^m.4, -11° 34'. This planetary is 44 by 26 seconds of arc in size, and of magnitude 8.4.

Beginners will also want to see NGC 7089 (M2), to be found at 21^h 30^m.9, -1° 03'. This is another globular cluster, a tight ball of stars 12 minutes in diameter, which glows at 7th magnitude. A 2-inch will show it well on a good night.

These three are the great well-known objects in the constellation, but the Skalnate Pleso *Atlas* shows others that tempt experienced seekers of deep-sky wonders. NGC 7184 is a spiral galaxy at 21^h 59^m.9, -21° 04', discovered by Sir William Herschel. Although only of about the 12th magnitude, its lenticular 5-by-1-minute shape is usually easily discerned.

At nearly the same declination as NGC

7184, but farther east, is the great planetary nebula of Aquarius, NGC 7293, 15 by 12 minutes of arc, situated at 22^h 27^m.0, -21° 06'. This was not seen by the early celestial explorers, for although it has a total magnitude of 6.5, its surface brightness is very low. It is a challenge to the observing amateur. The problem of measuring the distances of this and other planetaries was discussed by Otto Struve in the August issue, page 468, and NGC 7293 was pictured on page 208 in his article last March.

There are three interesting objects in the northwestern part of Aquarius. The globular cluster NGC 7492 can be picked up at 23^h 05^m.7, -15° 54'. Only 3.3 minutes in diameter and of the 12th magnitude, this is a difficult object, and its true nature is not obvious even in fair-sized telescopes.

Nestled in the midst of some naked-eye stars in NGC 7606, a spiral galaxy at 22^h 16^m.5, -8° 46', appearing as a narrow oval about six minutes long, its magnitude published as 11.5. Herschel, the first who saw it, described this nebula as easy, and it still is today.

Finally, the spiral galaxy NGC 7727, of magnitude 10.7, is located at 23^h 37^m.3, -12° 34'. It is an easy object in my 10-inch, shapeless and about two minutes in diameter.

WALTER SCOTT HOUSTON
Rte. 3, Manhattan, Kans.

OCCULTATION OF SATURN

On August 31st, the disappearance of Saturn behind the moon could be successfully observed from New Orleans, although the two objects were so low in the southwestern sky as to be among trees. The gradual approach of Saturn to the moon's dark edge was recorded photographically with a 3½-inch refractor on Plus-X film, by projection through a 40x eyepiece.

JOSEPH MICKEL
1307 Ridgeland Drive
Metairie, La.

Astronomy Study Aids



No. 736 Celestial-Terrestrial Globe. Clear plastic. Rotating terrestrial globe mounted inside star globe with unique star pointer. Globes show stars and constellations, countries and oceans, all in color. Demonstrates earth-sky-sun relationships and simplifies basic astronomy and geography.

No. 736-ST-12. 12" Celestial, 6" Terrestrial Globe, on cradle base. **\$42.50**

No. 736-ST-20. 20" Celestial, 6" Terrestrial Globe, on tripod base. **\$106.25**



No. 740-5 Deluxe Celestial Globe. Varicolored, shows all stars of the first six magnitudes, constellations, declination and right ascension circles, Magellanic Clouds and Milky Way. Important star clusters and nebulae indicated. Weighted base, movable meridian with horizon.


\$67.50

Science Associates

INSTRUMENTS/WEATHER
ASTRONOMY/TEACHING AIDS

Write for full catalogue.

194 Nassau Street, P.O. Box 216
Princeton, N. J.



SEPTEMBER AURORAE


DURING the great September naval maneuvers conducted in the North Atlantic by the combined fleets of the NATO powers, much difficulty was encountered in maintaining radio communications. This was a consequence of the intense solar activity during that period, which resulted in widespread and protracted disturbances of the earth's ionosphere and magnetic field. Teletype messages were garbled, and millions of Americans complained of the poor television reception that resulted from the sun's disturbed state.

In almost every part of the country, amateur astronomers observed through September a remarkable series of northern lights. The month was in its opening hours when Theodore L. Agos, of Worcester, Massachusetts, secured the pictures reproduced here, of a very brilliant display seen low in the northern sky from the summit of Mt. Wachusett.

He pointed his Super Ikonta B camera in the directions northwest, north-northwest, and north. By means of such star



groups as the Big Dipper, the pictures have been matched in azimuth, to make evident the pronounced changes in position and form of the auroral features in



Above and right: Three views of the fine northern lights of September 1st, photographed about 12:30 a.m. EST by T. L. Agos in central Massachusetts. He used an $f/2.8$ lens and Kodak Royal-X Pan 120 roll film. The nearly horizontal handle of the Big Dipper appears at the right in the second picture and the bowl is in the third picture.

Left: On September 3rd, about 3 o'clock in the morning, A. J. Morehouse at Battle Creek, Michigan, secured this one-minute exposure on Tri-X film at $f/4.5$. He pointed his Kodamatic camera 20 degrees up the north-western sky.



the intervals between the two-minute exposures.

Another reader of this magazine, Dr. J. R. Otoupalik, reports a spectacular aurora at Greeley, Colorado, two mornings later, about 3 a.m. Mountain standard time on September 3rd. From the northern quarter of the horizon, greenish-white rays extended 70 degrees up the sky, with the rising flashes that characterize a flaming aurora. A photograph of this same event, by A. J. Morehouse at Battle Creek, Michigan, is shown on this page.

Both of these displays were dwarfed by the widespread auroral storm on the evening of September 4th, which was extremely brilliant despite the presence in the sky of the moon only four days before its full phase. This display is perhaps to be classed with the greatest aurorae of the past few decades. From as far south as Batesville, Arkansas, J. R. Wright reports: "I first noticed the spectacle as a yellow-

green glow in the north at 8 p.m. Central standard time. By 9:40 p.m. the aurora reached the height of its activity. At that time there was a bright rayed arc in the north with rays that formed, moved toward a point south of the zenith, and faded. The dominating color of the forms was red, although the lower edge of the rayed arc was yellow-green. By 10 o'clock the rayed arc and rays had disappeared."

Other more northerly observers, favorably placed geographically, describe the spectacular corona. Kimball I. Jack, of Spokane, Washington, observed at Glacier Park in Montana, where northern lights covered three-fifths of the sky. He tells how rays like searchlight beams reached from the horizon practically to the zenith, converging near the constellation Lyra. But the region of convergence was dark and circular, appearing much like an open umbrella with a hole in the top. The coronal rays extended at least 15 degrees into the southern sky. Although remaining fixed in position, these features alternately faded and brightened.

Dale P. Cruikshank and R. J. Welch relate their impressions of the September 4th aurora, observed at Des Moines, Iowa, between 9:35 and 10:52 p.m. CST:

"It was the finest display seen this year. At one time, for about five to seven minutes, the aurora completely covered the entire sky, which was very clear. The gibbous moon (31 degrees above the south horizon) was engulfed, and streamers could be seen extending to the southern horizon between the trees."

These two observers noted the extreme rapidity of pulsations of the auroral forms; at 9:45 p.m. the northern sky was so bright that the 1st-magnitude star Capella could not be distinguished. Mr. Welch had already seen a twilight aurora that evening, marked by rapid pulsations.

Mrs. C. E. Fencken, at Hoopeston, Illinois, watched from 8:55 to 10:30 p.m. CST. Toward the display's climax two

very large and bright red patches formed in the east and west.

Other reports of September 4th have been received from C. E. Ott, Norwalk, Wisconsin; I. Alexeff, Madison, Wisconsin; T. V. Eynden, St. Bernard, Ohio; and R. R. Powell, Lawrence, Kansas.

Auroral activity continued at a high level throughout the month. Another bright display was observed only four days

The September 4th aurora was so bright at Pittsburgh that Walter A. Feibelman's photograph, at 11:03 p.m. EST, shows the rayed structure very distinctly, despite the interference by clouds and bright moonlight. Mr. Feibelman noted that maximum auroral intensity occurred near 10:50 or 10:55 p.m. Some of his pictures record rayed arcs with the parallel rays very thin and numerous. As seen elsewhere under better sky conditions, this display was among the most extraordinary in years.



after full moon by Walter A. Feibelman, who makes systematic photographic records at Pittsburgh, Pennsylvania, in cooperation with the IGY Auroral Data Center at Cornell University. He commented on the wide extent of this aurora, reaching far into the west and southwest. A bright red spot appeared near the close of the display.

A noteworthy auroral storm on the night of September 22-23, widely seen over the United States, is discussed by James E. McDonald in his article on page 14. R. T. Morehead, at Des Moines, Iowa, describes this aurora as consisting of rays converging on the zenith, some extending down nearly to the southern horizon. Almost half the sky was brilliant red at the height of the display, a few rays appearing almost blood red; in the northern sky, however, the forms were mainly greenish white.

On the West Coast, at Vancouver, British Columbia, Terry Taylor watched the peak of the September 22nd display at 8:50 p.m. Pacific standard time. He writes, "The corona had grown into a spectacular red-and-white whirlpool of milky light, with a clear spot in the center. Bright rays covered over 75 per cent of the sky, nearly obliterating the 2nd-magnitude stars in Cassiopeia. In the densest

regions of the corona even Vega and Deneb looked inconspicuous."

The next night, September 23-24, all the stages of development and decay of a lesser aurora at Pittsburgh were photographed by Mr. Feibelman within less than an hour.

Chris Schell saw two aurorae during the month, one on the morning of September 2nd from Ithaca, New York, the other on


September 29th from Pittsfield, Massachusetts. The latter was mainly a drapery formation, with strong red patches.

The northern lights reported here can at best comprise only a small part of the total auroral activity during this memorable month, for they are merely a sampling from a limited geographical area. Only when systematic records of the IGY's world-wide auroral program have been analyzed will the over-all story be learned.

COMET MRKOS AND SOLAR ACTIVITY

Recent photographs of Comet Mrkos, according to astronomers at Yerkes Observatory, show the occurrence in its tail of irregular, changing patches of light. These patches are composed of ionized gases, whose luminosity is caused by jets of fast-moving electrons, protons, and alpha particles ejected from the sun. Similar streams of solar corpuscular radiation that collide with the earth are responsible for the bright northern lights observed during the present peak in the 11-year cycle of solar activity.

Comet Mrkos is the object that was so conspicuous during August, but its rapid recession from the sun has now carried it into the southern sky, where it is very faint.



UNITRON Altazimuth Refractors: left to right, 1.6" (with star diagonal), 2.4" (with erecting prism system), 3" (with DUETRON double eyepiece), and 4" (with astro-camera and with UNIHUX on the shelf).



UNITRON Equatorial Refractors: left to right, 2.4" (with astro-camera), 3" Photo-Equatorial (with sun screen), and 4".

Does Santa know you want a UNITRON for Xmas?

MANY Models To Choose From!

1.6" ALTAZIMUTH (\$7.50 Down)	\$75
with eyepieces for 78x, 56x, 39x	
2.4" ALTAZIMUTH (\$12.50 Down)	\$125
with eyepieces for 100x, 72x, 50x, 35x	
2.4" EQUATORIAL (\$22.50 Down)	\$225
with eyepieces for 129x, 100x, 72x, 50x, 35x	
3" ALTAZIMUTH (\$26.50 Down)	\$265
with eyepieces for 171x, 131x, 96x, 67x, 48x	
3" EQUATORIAL (\$43.50 Down)	\$435
with eyepieces for 200x, 131x, 96x, 67x, 48x	
3" PHOTO-EQUATORIAL (\$55.00 Down)	\$550
with eyepieces for 200x, 171x, 131x, 96x, 67x, 48x	
4" ALTAZIMUTH (\$46.50 Down) with	\$465
eyepieces for 250x, 214x, 167x, 120x, 83x, 60x	
4" EQUATORIAL (\$78.50 Down) with	\$785
eyepieces for 250x, 214x, 167x, 120x, 83x, 60x, 38x	
4" PHOTO-EQUATORIAL (\$89.00 Down) with	\$890
eyepieces for 250x, 214x, 167x, 120x, 83x, 60x, 38x	
4" EQUATORIAL with clock drive	\$985
(\$98.50 Down), new Model 160V	
4" EQUATORIAL with clock drive and	\$1075
metal pier (\$107.50 Down), new Model 166V	
4" PHOTO-EQUATORIAL with clock drive and	\$1175
astro-camera, (\$117.50 Down), with eyepieces	
for 250x, 214x, 167x, 120x, 83x, 60x, 38x, 25x	
4" PHOTO-EQUATORIAL with clock drive, pier,	\$1280
astro-camera (\$128.00 Down), eyepieces for 375x,	
300x, 250x, 214x, 167x, 120x, 83x, 60x, 38x, 25x	

Higher- and lower-power eyepieces available for all models. Prices include basic accessories, tripod and mounting, fitted wooden cabinets, and operating instructions. Additional accessories available to add further to your observing pleasure.

NOW is definitely the time to reserve a UNITRON for that very special person on your Christmas list who perhaps may be none other than yourself. Precision craftsmanship cannot be hurried and the number of UNITRONS which Santa will have available is necessarily limited. To avoid disappointment, order your UNITRON now. Full payment or the down payment may be made at the time you wish to have the telescope delivered. Remember, there is no substitute for a UNITRON.

• LENSES are FULLY CORRECTED for spherical and chromatic aberration and are COATED for maximum brilliance and clarity of image. Optics especially designed for astronomical observation. AIR-SPACED OBJECTIVES.

• REFRACTOR type of design duplicates the performance of larger telescopes of other types. No mirrored surfaces to become oxidized. Superior definition to the very edge of the field.

• EYEPIECES of the HIGHEST QUALITY: Orthoscopic, Achromatized Symmetrical, Kellner, Huygens. Three to ten eyepieces included with each instrument as standard equipment.

• FINEST MATERIALS throughout. DURALUMINUM TUBE. Moving parts of BRASS carefully machined to close tolerances, and finished in CHROMIUM. No war-surplus components used.

• MODERN DESIGN based on time-tested engineering principles. HANDSOME APPEARANCE to which no illustration can do justice.

• EQUATORIAL MODELS have slow-motion controls for both declination and right ascension as well as rapid-motion controls. Sturdy TRIPOD.

• ALTAZIMUTH MODELS have slow-motion controls for both altitude and azimuth as well as clamps for both co-ordinates. Sturdy TRIPOD.

• VIEW FINDER with crosshair eyepiece.

• RACK-AND-PINION FOCUSING.

• Choice of UNIHEX Rotary Eyepiece Selector or STAR DIAGONAL and ERECTING PRISM for TERRESTRIAL OBSERVATION. The same complete range of terrestrial magnifications as for celestial observation.

HOW TO ORDER

Send check or money order in full or use our Easy Payment Plan, as described below. Shipments made express collect. Send 20% deposit for C.O.D. shipment. UNITRON instruments are fully guaranteed for quality, workmanship, and performance, and must meet with your approval or your money back.

Use Our EASY PAYMENT PLAN

UNITRON's popular Easy Payment Plan is a convenient and economical way to buy your UNITRON Refractor when you do not want to disturb your savings or when you haven't the total cost of the telescope immediately available. The down payment required is 10%. The balance due is payable over a 12-month period, and there is a 6% carrying charge on the unpaid balance. Your first payment is not due until 30 days after you receive the instrument, and if you should want to pay the entire balance due at that time, the carrying charge is canceled.

There is no "red tape" when you order—you merely fill in the UNITRON Easy Payment Order Form, return it together with the required down payment, and the model of your choice is shipped to you at once.

Do not hesitate to use this plan merely because you have never before purchased anything on time payments. The procedure is practically painless and is an ideal means of enjoying the use of your UNITRON while paying for it.

Telescope Accessories by UNITRON

UNITRON ACCESSORIES include objective lenses, eyepieces, rack-and-pinion focusing mechanisms, sun projecting screens, guide telescopes, the UNICLAMP camera brackets, and many other components in addition to those listed on this page. All are described, illustrated, and priced in the new UNITRON Catalog. Since each component is used as standard equipment in UNITRON Refractors themselves, you are assured of obtaining the finest quality.

ASTRO-CAMERA 220: Complete with air-operated curtain shutter, giving speeds of 1/10 to 1/90 second, bulb and time, shutter release, ground-glass back, 3 double plateholders for 3 1/4" x 4 1/4" plates or cut film, a 30-mm. f.l. eyepiece, extension tubes and clamps, and a fitted wooden cabinet. Model A is designed to fit the UNITRON rack and pinion and is for UNITRONS only. Model B fits 1 1/4" eyepiece holders. A or B:

Only \$69.50 postpaid

DUETRON Double Eyepiece: With DUETRON, two observers may use the telescope simultaneously and with equal comfort. The eyepieces used need not necessarily be of the same magnification. DUETRON is a real boon for father-and-son observing teams. With DUETRON, more observers may be accommodated at star parties, and the advanced members may provide valuable instruction to the beginners. The tube of DUETRON Model A is interchangeable with the customary UNITRON drawtube and may be used with all models with the exception of the 1.6" Refractor. DUETRON Model B fits 1 1/4" eyepiece holders. Complete with special clamping device and cabinet. A or B:

Only \$23.50 postpaid

1. VIEW FINDER (As shown on UNITRON 2.4" Equatorial): 23.5-mm. (.93") achromatic objective, 6x eyepiece with crosshairs. Chromed brass tube. Mounting brackets with centering screws.

Only \$8.50 postpaid

2. VIEW FINDER (As shown on UNITRON 3" Altazimuth): 30-mm. (1.2") coated achromatic objective and 8x eyepiece with crosshairs. Other details as in View Finder 3.

Only \$10.75 postpaid

3. VIEW FINDER (As shown on UNITRON 4" Refractors): 42-mm. (1.6") coated achromatic air-spaced objective. 10x eyepiece with crosshairs. Duraluminum tube finished in white enamel. Dewcap. Furnished with mounting brackets with centering screws for collimation. This finder also makes an excellent hand telescope for spectacular wide-field views of the sky.

Only \$18.00 postpaid

UNIHEX ROTARY EYEPIECE SELECTOR: The old-fashioned method of fumbling with eyepieces in the dark has been outmoded by UNIHEX, UNITRON's new Rotary Eyepiece Selector. With UNIHEX, you always have 6 magnifications ready at your fingertips. To change power, merely rotate a new eyepiece into position while the object stays centered and in focus in the field of view. Model A is designed to fit the UNITRON rack and pinion and is for UNITRONS only. Model B fits 1 1/4" eyepiece holders. Complete with special clamping device and cabinet. A or B:

Only \$24.75 postpaid

Get UNITRON'S FREE

Observer's Guide and Catalog

on Astronomical Telescopes.

This valuable 38-page book is yours for the asking!

With the artificial satellite and space travel almost a reality, astronomy has become today's fastest growing hobby. UNITRON'S new handbook contains full-page illustrated articles on astronomy, observing, telescopes and accessories. Of interest to beginner and advanced amateurs alike.

Contents include —

- Observing the sun, moon, planets and wonders of the sky
- Constellation map
- Hints for observers
- Glossary of telescope terms
- How to choose a telescope
- Amateur clubs and research programs



UNITRON
Instrument Division
of UNITED SCIENTIFIC CO.

204-6 MILK STREET • BOSTON 9, MASS.

Please rush to me, free of charge, UNITRON'S new Observer's Guide and Telescope Catalog.

Name.....
Street.....
City.....State.....

UNITRON

INSTRUMENT DIVISION OF
UNITED SCIENTIFIC COMPANY

204-206 MILK STREET • BOSTON 9, MASSACHUSETTS

ENTERTAINMENTS, PUZZLES . . .

T167. THE COMPLETE NONSENSE OF EDWARD LEAR. All original text, illustrations of all nonsense books Lear ever wrote. 5 books in 1. "Dong with a Luminous Nose," "Owl and Pussycat" — hundreds of others. 546 illustrations. 232 pp. \$1.00

T1. FLATLAND. E. A. Abbott. Fictionalized life in 2 dimensions, offers techniques for visualizing multidimensional forms, etc., clearer than most texts. "Delightful," *Science News Letter*. 128 pp. \$1.00

T97. CRYPTANALYSIS. H. F. Gaines. Best intermediate text for serious student; modern techniques, much not usually known except to experts. 236 pp. \$1.95

T362. LEARN CHESS FROM THE MASTERS. F. Reinfield. Play 10 games against masters and grade every move you make! 152 pp. \$1.00

T162. MATHEMATICAL RECREATIONS. M. Kraitchik. Moderately advanced problems based on theory of numbers, topology, etc. Unequalled for easily assimilated insights. 330 pp. \$1.75

T337. PUZZLE QUIZ STUNT FUN. Jerome Meyer. Mathematical puzzles, deductions, observations. 238 in all. Entirely new. 256 pp. \$1.00

T344. NEW WORD PUZZLES. G. L. Kaufman. 100 entirely new puzzles for enthusiasts of word games: chesswords, designonyms, addie-letter anagrams, lapwords, acrostigrams, etc. 196 figures. 128 pp. \$1.00

T110. MATHEMAGIC, MAGIC PUZZLES, GAMES WITH NUMBERS. R. V. Heath. One of most famous books in recreational math. 128 pp. \$1.00

T384. HOUDINI ON MAGIC. Houdini's own account of magic tricks, picking locks, exposes of spiritualism, how stage magic works, etc. Incredible feats of greatest magician of modern times. 155 photos. 295 pp. \$1.00



T198. MATHEMATICAL PUZZLES FOR BEGINNERS AND ENTHUSIASTS. G. Mott-Smith. 188 mathematical puzzles of most delightful sort, detailed solutions. 248 pp. \$1.00

T143. BOOK OF MODERN PUZZLES. G. L. Kaufman. More than 150 word and logic puzzles, unusual types, highly original; no old chestnuts. 192 pp. \$1.00

SPECIAL ANNOUNCEMENT!

S401-3. LES METHODES NOUVELLES DE LA MECHANIQUE CELESTE. Henri Poincaré. Complete text in French, of Poincaré's great monograph. Presents first use of integral invariants, important applications of linear differential equations, investigation of periodic orbits, other important celestial problems. Approaches 3-body problem from many different ways. Lunar motion, satellites of Jupiter. Thorough, complete coverage. Total of 1,280 pp. 3-volume set. \$7.50

Can you explain this?



Rearrange the parts of the top figure into the bottom one, and you lose one square of area. Mysterious, isn't it?

Let MARTIN GARDNER, well-known puzzle editor, explain this and hundreds of other elements of mathematical magic!


MATHEMATICS, MAGIC AND MYSTERY

by Martin Gardner

How do geometrical vanishing tricks work? Why are card tricks possible? How can magicians do astonishing feats of mathematics mentally? As a rule, we accept these tricks as "magic," and do not recognize that they are demonstrations of probability, theory of numbers, topology, and other mathematics.

This is the first book-length study of this fascinating branch of mathematics. Written in the same delightful manner as Gardner's articles in *Scientific American*, it summarizes all previous work in this field. It offers scores of new tricks, insights, puzzles, and aligns many effects with law for the first time. It discusses tricks with cards, coins, topological tricks, geometric vanishing effects, and the like. You will see how Sam Loyd could make a Chinaman vanish, how a Curry square "proves" that the whole is more than its part.

T335. 135 illustrations. xii + 174 pp. \$1.00



NOW only

\$2.45!

FOUNDATIONS OF PHYSICS

by R. B. Lindsay and H. Margenau

PARTIAL CONTENTS

I. MEANING OF PHYSICAL THEORY. Data, experiment, measurement, symbolism, laws, theory, abstraction, boundary conditions, integral, differential equations. — II. SPACE, TIME. Psychology vs. physics, geometry. — III. FOUNDATIONS, MECHANICS. Kinematics, position, displacement, velocity, acceleration, dynamical concepts, Newton, mass, force, motion, d'Alembert, Gauss, Hertzian mechanics, energy, canonical equations of Hamilton. — IV. PROBABILITY, APPLICATION. Probability aggregates, Bernoulli's problem, emission of rays, Laplace's formula, density fluctuations. — V. STATISTICAL POINT OF VIEW. Dynamical, statistical theories, electric current, thermodynamics, statistical mechanics, Gibbs, Darwin, Fowler. — VI. PHYSICS OF CONTINUUA. Continuum, scalar, vector fields, deformable media, tensor fields, electromagnetic fields. — VII. ELECTRON THEORY, SPECIAL RELATIVITY. Invariance of field equations, special theory. — VIII. GENERAL THEORY. Ideas, motion of particle in gravitational field. — IX. QUANTUM MECHANICS. Uncertainty, Hermitean operators, Schrödinger's equation, eigenvalues, matrices, spin, quantum statistics, Dirac. — X. CAUSALITY.

S413. INTRODUCTION TO STUDY OF STELLAR STRUCTURES. S. Chandrasekhar. Unabridged republication of modern classic, costing up to \$20 on out-of-print market. 509 pp. \$2.75

S366. APPLIED OPTICS AND OPTICAL DESIGN. A. Conrady. Full unabridged inexpensive reissue of standard modern treatise. 528 pp. \$2.95

T203. ANIMALS IN MOTION. Eadweard Muybridge. Largest collection (over 4,000 photos) ever made from famous Muybridge collection of high-speed stopped-action photos of animals in motion. Horses, cats, lions, guinea pigs, camels, sloths, elephants, 27 other animals, birds, 123 different types of action. Invaluable for biologists, zoologists, naturalists. New introduction by L. S. Brown, American Museum of Natural History. Over 410 pp. 7 7/8" x 10 3/8". Deluxe edition. Clothbound \$10.00

T295. HOW TO CALCULATE QUICKLY. H. Sticker. Tried and true method, over 9,000 problems, to develop your number sense. Techniques not taught in school: division, multiplication, etc. 256 pp. \$1.00

S109. VECTOR AND TENSOR ANALYSIS. G. E. Hay. Start with simple definitions, finish understanding vectors and tensors. 201 pp. \$1.75

S168. NUMERICAL SOLUTIONS OF DIFFERENTIAL EQUATIONS. Levy and Baggott. Unequalled time saver. Graphic, numerical methods. "Should be in hands of all in research in applied math." *Nature*. 246 pp. \$1.75

S133. TABLES OF FUNCTIONS, with formulae, curves. E. Jahnke and F. Emde. Most comprehensive 1-volume collection. 394 pp. \$2.00

S164. INTRODUCTION TO SYMBOLIC LOGIC. Susanne Langer. Even if you never finished high school algebra you can learn modern Russell-Whitehead logic. 368 pp. \$1.75

S27. NON-EUCLIDEAN GEOMETRY. R. Bonola. Standard coverage, history, exposition, with complete texts of Bolyai and Lobachevski included. 180 diagrams. 431 pp. \$1.95

S182. GEOMETRY OF FOUR DIMENSIONS. H. P. Manning. Unique in English, simple, easily followed account. 76 figures. 357 pp. \$1.95

S292-3. THEORY OF SOUND. Lord Rayleigh. Solutions for vibrating systems, hints on nonlinearities. Harmonic functions, theory of resonators, etc. Formerly \$12. Total 1,041 pp. 2-volume set. \$3.90

S81. PRINCIPLE OF RELATIVITY. Einstein, Lorentz, Weyl, and Minkowski. 11 papers founding relativity. Unabridged translations. "Indispensable," *Nature*. 224 pp. \$1.65

S193. HIGHER MATHEMATICS FOR STUDENTS OF CHEMISTRY AND PHYSICS. J. Mellor. Class favorite, unequalled review, practical rather than theoretical on lab situations; integral, differential calculus, theory of errors, etc. 662 pp. \$2.00

S263. PARTIAL DIFFERENTIAL EQUATIONS OF MATHEMATICAL PHYSICS. A. G. Webster. Special methods, insights, standard coverage. 448 pp. \$1.98

"Thorough and yet not over-detailed . . . unreservedly recommended," *NATURE* [London].

A bridge between semipopular works and technical treatises, this book analyzes foundation ideas and methods of modern theoretical physics. It is not a textbook but a discussion of methods of physical description and construction of theory with full cognizance of the logic of science, physical data and their measurement, construction of experiment, and symbolic representation.

Intended for the intermediate or advanced reader who has some background in the sciences and some knowledge of calculus, it covers both classical and modern physics, contrasting their positions, and telling why transitions were made from classical to modern. It gives meaning to the bare data and tools of physics.

Reviews of the first edition: "A most unusual book . . . an interestingly fresh approach . . . will help [the reader] to discover that there is much more to his subject than is involved in adroit manipulation of differential equations." *E. U. Condon, Review of Scientific Instruments*. "At once thorough and yet not over-detailed; it may be unreservedly recommended," *Nature* [London].

S377. Index. Reading lists. 35 illustrations. xi + 537 pp. \$2.45

T394. FADS AND FALLACIES IN THE NAME OF SCIENCE. Martin Gardner. Enlarged revised edition of *In the Name of Science*, the standard coverage of the cults, fallacies, and curious systems that have recently masqueraded as science. It ranges from astronomy to theoretical physics, chemistry, geology, medicine, psychology, philosophy, history, scientific method. It is not a collection of anecdotes, but a fair, reasoned appraisal of eccentric theory. Flying saucers, Atlantis, general semantics, dianetics, orgone energy, hollow earthism, Bridey Murphy—dozens of others are all treated. "Every scientist, for the preservation of his own good sense, should read this book," *Quarterly Review of Biology*. "Hilariously entertaining," *San Francisco Chronicle*. x + 365 pp. \$1.50

S267. SPACE TIME MATTER. H. Weyl. "Standard treatise on the general theory of relativity," *Nature*. 348 pp. \$1.75

S361. THERMODYNAMICS. Enrico Fermi. Clear, elementary. "May be recommended as clear, very accurate introduction," *Journal of Physical Chemistry*. 170 pp. \$1.75

S298. FAMOUS PROBLEMS OF ELEMENTARY GEOMETRY. F. Klein. Squaring circle, doubling cube, pi, in terms of rigorous modern math. 103 pp. \$1.00

S342. INTRODUCTION TO THEORY OF NUMBERS. L. Dickson. Nontechnical, wide coverage. 239 problems. 191 pp. \$1.65

T371. TRIGONOMETRY REFRESHER FOR TECHNICAL MEN. A. Klaf. 913 detailed questions, answers, brush-up for you. 1,738 problems. 436 pp. \$2.00

T370. CALCULUS REFRESHER FOR TECHNICAL MEN. A. Klaf. 756 detailed questions, answers, in unique practical refresher. 566 problems. 436 pp. \$2.00

All books standard 5 7/8" x 8" and PAPERBOUND unless otherwise indicated.

Dept. 119, Dover Publications, Inc.
920 Broadway, New York 10, N. Y.

Please send me the books encircled below.
(Please print)

Name

Address

City Zone State

I am enclosing \$..... in full payment. (Payment in full must accompany all orders except those from libraries or public institutions, who may be billed.) Please add 10c postage per book on orders less than \$5.00. Dover pays postage on orders of \$5.00 or more.

T1	S133	S182	S292-3	T344	S377
S27	T143	S193	T295	S361	T384
S81	T162	T198	S298	T362	T394
T97	S164	T203	T335	S366	S401-3
S109	T167	S263	T337	T370	S413
T110	S168	S267	S342	T371	X773

Satisfaction guaranteed! All books may be returned within 10 days for full cash refund if they are unusable for any reason whatever.

BOOKS AND THE SKY

LIGHT SCATTERING BY SMALL PARTICLES

H. C. van de Hulst. John Wiley and Sons, New York, 1957. 470 pages. \$12.00.

ALMOST ALL information used by astronomers comes to us in the form of electromagnetic radiation, or light. For many purposes, it is sufficient to think of the light as traveling unimpeded through empty space from the source to the observer.

Many concepts of astronomy are based on this simplification. For example, in positional astronomy we assume that the light has traveled in a straight line from the star or planet. In calculating absolute magnitudes, or in determining distance through absolute magnitude effects, we assume that the intensity of the light falls off as the inverse square of the distance.

In most cases, however, when we derive physical information about astronomical bodies or the interstellar medium, we have to take into account the interaction between the radiation and the material it is traversing. (Indeed, we must also do this in designing telescope optics for examining the radiation.) When Fraunhofer absorption lines are produced by a stellar atmosphere, the radiant energy does not come to our instruments directly from the source, but is reflected, absorbed and re-emitted many times in the course of its passage. Such interactions form much of the subject matter of astronomy and physics.

Studies of colloidal solutions and aerosols in chemistry, scattering of X-rays in physics, viruses and blood corpuscles for biology, haze, fog, and rain droplets in meteorology, all make use of information derived through the scattering of light by small particles. The work done for one particular application is frequently of use in a completely different discipline.

It is often more important to consider the effect of single small particles one at a time than that of the transmitting medium as a whole. In astronomy this is true for investigations of interstellar and interplanetary dust, as well as of the atmospheres of other planets. For this reason, and in order to make a book of practical size, the author has severely limited its scope.

Processes producing a change in wave length of the light are ruled out. Dr. van de Hulst considers only scattering by well-separated particles, where the interaction can be considered to involve only one particle at a time. He does not discuss multiple scattering, in which the light is scattered again by a second particle.

Within these limitations, perhaps because of them, the book will be invaluable to research workers in many fields, but it is not particularly suitable as a textbook, even for an advanced course in optics. Its

primary value will be as a reference providing in one place a logical and concise summary of scattering theory. Its mathematical nature makes the book of little use to most amateur astronomers.

Dr. van de Hulst discusses first the basic theory of scattering. For particles about the size of the wave length of the light involved, actual calculations of scattering are very tedious, and it is not yet possible to compute the scattering by arbitrarily shaped particles for all wave lengths and polarizations. The second and largest section of the book deals with the theories of scattering by many special types of particles. A final section gives some typical applications to chemistry, meteorology, and astronomy, in a brief form sufficient to indicate to the research worker what parts of the scattering theory may be applicable to his particular problem. One of the most important of these in astronomy is the effect of interstellar grains in producing absorption and polarization of light that passes through the space within the galaxy.

The author says that his needs in astrophysical research primarily led him to write this review of the widely dispersed literature on scattering. Nonetheless much of his own work appears in many parts of the volume, indicating his contributions to gaps in the available theory.

ROBERT FLEISCHER
Rensselaer Polytechnic Institute

DISCOVERIES AND OPINIONS OF GALILEO

Stillman Drake, translator. Doubleday Anchor Books, Garden City, New York, 1957. 302 pages. \$1.25, paper bound.

FOUR selections from Galileo's works are translated here. An introduction to each piece gives the biographical background and some details on the circumstances that led to its writing.

The first two selections, "The Starry Messenger" and "Letters on Sunspots," are particularly interesting from the astronomical point of view. The former especially conveys the excitement, as well as the brilliance, of Galileo's discoveries on first looking through a telescope. We are reminded that things now taken for granted, such as the irregularity of the lunar surface and the revolution of the satellites about Jupiter, were novel concepts 350 years ago.

In reading these works this reviewer was once more impressed by the genius of this man, who not only saw and described so many things new in his day, but usually interpreted correctly what he saw. This understanding of natural phenomena is particularly impressive in view of the considerable opposition to his ideas.

We may take as an example the case of

Announcing...

A STEP
BY STEP
HOW-TO-DO-IT
BOOK

HOW TO MAKE A TELESCOPE

By JEAN TEXEREAU

Translated and Adapted from the French by Allen Strickler. With forewords by André Couder and Albert G. Ingalls.

When the original French version of this famous book was published, Albert Ingalls, the well-known editor of the American classic "Amateur Telescope Making," said:

"... if translated into English it would make a splash in amateur telescope making circles."

In his foreword to the book, Mr. Ingalls states, "It is an especially well organized book which sticks entirely to the sequence of logic... the entire discussion pertaining to each stage of the work is completed before the book proceeds to the next stage... you will not get lost on the way..." As the editor of books on the same subject as this, I am happy to move over on the optical bench to make room for Jean Texereau and his book."

Jean Texereau started as an amateur and is now technical associate of the Optical Laboratory of the Paris Observatory and secretary of the Instrument Group of the Astronomical Society of France. His translator, Allen Strickler, is associated with Beckman Instruments Company.

CONTENTS:

1. Basic Principles and a Proposed Telescope.
2. Making the Large Mirror.
3. The Plane Diagonal Mirror.
4. The Eye Piece.
5. Mechanical Structure.
6. The Altazimuth Mounting.
7. Accessories, Mirror Coating, Adjustments.
8. The Telescope in Use.
9. Photography with the Amateur Telescope.
- Appendix.
- Sources of Telescope Making Supplies.
- Index.

1957. 208 pp., 86 ill., 7 tables. \$3.50

INTERSCIENCE PUBLISHERS, INC.
250 Fifth Ave., New York 1, N.Y.

Gentlemen: Please send me () copy of **TEXEREAU** \$3.50

() Check Attached
(For N.Y. City delivery add 3% Sales Tax)

Name _____
(PLEASE PRINT)

Address _____

City _____ Zone _____ State _____

☐ I would prefer to receive a copy of the descriptive folder before placing my order.

CHRISTMAS SHOPPING?

Why not GROUP SHOP among our specials?

Group No. 1

<i>The Sun</i> , G. Abetti	\$12.00
<i>The Moon</i> , Wilkins and Moore	12.00
<i>Physics of the Planet Mars</i> , G. de Vaucouleurs	10.00
A \$34.00 value for only	\$32.00

Group No. 2

<i>The Milky Way</i> , Bok and Bok	\$5.50
<i>Between the Planets</i> , F. Watson	5.00
<i>Earth, Moon, and Planets</i> , F. Whipple	4.75
<i>The Story of Variable Stars</i> , Campbell and Jaccia	3.50
An \$18.75 value for only	\$17.25

Group No. 3

<i>The Expansion of the Universe</i> , P. Couderc	\$6.00
<i>The Sun and Its Influence</i> , M. Ellison	4.50
<i>From Atoms to Stars</i> , M. Davidson	3.75
<i>Introducing Astronomy</i> , J. Sidgwick	3.50
A \$17.75 value for only	\$16.35

Group No. 4

<i>Frontier to Space</i> , E. Burgess	\$4.50
<i>Rocket Propulsion</i> , E. Burgess	4.50
<i>Guided Missiles in War and Peace</i> , N. Parson, Jr.	3.50
A \$12.50 value for only	\$11.75

NEW: *Discovery of the Universe*,
G. de Vaucouleurs \$6.00

As usual we will gift wrap your purchase on request, and enclose your gift card when sent with your order.

Write for our complete circulars on books, slides, charts, and other astronomical literature.

ASTRONOMY CHARTED

33 Winfield St., Worcester 10, Mass., U.S.A.

Phone: PL 5-6992

the interpretation of sunspots. After careful observations Galileo decided that they were very close to, if not actually on, the solar surface. Learned opinion on the whole was opposed to this, for it implied that there were blemishes on the sun's surface, and that it was not a pure ball of fire, as believed by the Aristotelians.

In the last two essays of the book, Galileo goes into the matter of scientific truth founded on observation and logical deduction, as opposed to authority (particularly Aristotle) and faith. He draws a strong distinction between matters that can be investigated scientifically, and those in the domain of theologians. Such a restricted viewpoint was necessary for survival in Renaissance Europe, but even today many scientific philosophers admit limitations to the methods of scientific thought.

Mr. Drake is to be congratulated for making these works of Galileo available to the general reader. The introductory notes give the background necessary to a fuller understanding of the selections, by describing the circumstances under which they were written. In several cases Mr. Drake points out instances where later work indicated that Galileo had erred. Somewhat more material of this kind might have been included for the benefit of the reader who knows nothing of astronomy.

The selections contained in this volume

are all from Galileo's early period, 1610 to 1623. His famous *Dialogue on the Great World Systems* (1638) is not included, but is currently in print.

A lack of knowledge of the original Italian versions of these works prohibits the reviewer from passing judgment on the exactness of the translation, but it reads easily and smoothly. The style is clearly modern, which makes for better reading.

ELSKE V. P. SMITH
Sacramento Peak Observatory

THE LIFE OF ARTHUR STANLEY EDDINGTON

A. Vibert Douglas. Thomas Nelson and Sons, New York, 1957. 207 pages. \$6.25.

WRITING the biography of a many-sided genius would have been difficult even under the best circumstances. In Sir Arthur Eddington's case the task is made more difficult because of his shy and retiring nature. As one whose interest in astronomy was kindled by the writings of Eddington, the reviewer had been looking forward to reading this work by Professor Douglas, who was a student and associate of the famous astrophysicist. She has done a commendable job, and fellow scientists will be indebted to her for providing an integrated view of the activities of a truly great scientist.

In the first two chapters are traced as far as is possible the early influences in Eddington's life that shaped his later interests. There is a vivid account of Eddington as a happy young man with prodigious intellectual curiosity and the ability to follow up this curiosity in any field he liked. That circumstances were favorable for his activities is both a commentary on his attitude to life and on the state of the world at that time.

After spending seven fruitful years as chief assistant at Greenwich, Eddington returned to Cambridge in 1913 as Plumian Professor. He was to remain in the same position until his death in 1944. It is sometimes forgotten that Eddington, who is best remembered for his theoretical work, during his years at Greenwich had "to observe at nights a good deal in order to understand the instruments thoroughly."

The author gives a fairly general but interesting account of Eddington's work, beginning with his early studies of star motions, through stellar structure, relativity, and cosmology. In every one of these fields he had the uncanny knack of selecting the most significant problems and creating whole new branches of astrophysics. There is also a lively narrative of his controversies with his fellow astronomers, especially Sir James Jeans. In these controversies we see the growing pains of astrophysics as a subject with its own rules of the game.

Apart from his astronomical researches, Sir Arthur was very active both in the

SKYSCOPE With standard 60-power eyepiece \$29.75

The full 3½-inch diameter reflecting-type astronomical telescope that even the telescope makers talk about.

It has been sold for more than 17 years and now is on display in at least two U. S. planetaria. It will show mountains and craters on the moon, Saturn's rings, Jupiter's four moons and the planet's markings, and close double stars, with guaranteed observatory clearness. Skyscope enjoys world-wide distribution.


Every instrument, with its ¼-wave, aluminized mirror, is individually tested before being packed for shipment. We suggest that before buying you inquire at almost any local astronomy society about the efficiency of Skyscope. 100% American-made.

We invite your attention to our free and straightforward descriptive brochure which also shows a photograph of the individual parts used.

125-power and 35-power extra eyepieces	\$5.15 each
6-power finder, with brackets	\$7.50
Holder for extra eyepieces	\$1.00

THE SKYSCOPE CO., INC.

475-s Fifth Avenue, New York 17, N. Y.



The latest

Spitz MODEL A-1 PLANETARIUM

has been shipped to
**University
of Nebraska
State Museum**
Lincoln, Nebraska



**Spitz Laboratories
Inc.**

YORKLYN, DELAWARE
Phone: CEder 9-5212

popularization of science and the development of a philosophy of science. In later life, one of the criticisms by his less broad-minded colleagues was that he mixed mysticism and science instead of using his brilliant analytical talents for scientific study only. As the book indicates, "one of the most powerful factors in the formation of Eddington's intellectual outlook and spiritual perceptivity was the Quaker atmosphere of his home."

In a chapter titled "Concerning the Unseen World," Miss Douglas gives a lucid outline of Eddington's personal philosophy. In these days of unquestioned obedience to specialization this outlook is like a breath of fresh air. One gets a glimpse of the essential morality and the humility of the man who could remark at the end of a classic on stellar structure, "Somewhere in the present tangle of evolution and sources of energy I have been misled; and my guidance of the reader must terminate with the admission that I have lost my way." Eddington did not isolate his scientific work from his philosophy of life, and he felt a deep moral responsibility for all his actions. There may have been more brilliant scientists, but one doubts whether there have been many endowed with his wisdom.

The penultimate chapter is devoted to the most controversial of all Professor Eddington's works. Most of his last years

were occupied in building up an elaborate theory of the microscopic and macroscopic structure of the universe. The final portions of this study, entitled *Fundamental Theory*, were published posthumously. The whole work is so abstruse that even a well-known cosmologist remarked, "The analysis . . . is so very complex and based on such extremely difficult arguments that most scientists found themselves unable to examine the theory in detail."

No review of Miss Douglas' book can be complete without commenting on the literary excellence of the biography of a man who was noted for his facility with the pen and the imagery of his descriptions of extremely difficult concepts. Studded with generous quotations from Eddington's writings, this biography will provide fascinating reading for both the professional and amateur astronomer.

T. K. MENON
Harvard College Observatory


OPTICS: THE SCIENCE OF VISION

Vasco Ronchi. New York University Press, New York, 1957. 360 pages. \$10.00.

THE DIRECTOR of the National Institute of Optics, Arcetri, Italy, who is well known as the inventor of the Ronchi test for optical systems, has not given us here a textbook in optics, as the title might imply. Rather this is a pioneering work, pleading for a science of vision to bring together those aspects of physics, physiology, and psychology that deal with how we see, both with and without the aid of telescopes and microscopes.

Such a science would not be solely a chapter in physics, where the eye is generally treated as something placed at the eyepiece of an instrument. Nor would this new field be merely a part of physiology, where optical instruments are only devices placed in front of that complex organ, the eye. The science of vision would have a broader outlook than psychology, where the optical instrument and eye are both simply media for transmission of stimuli to certain parts of the brain. There, in a little-understood and seemingly magical manner, arises a vision of a rose garden, a speeding car, or a galaxy in Andromeda. The new science would aim at the totality of what these three disciplines have to say on the subject.

In short, Ronchi would define optics as the science of vision, not merely the study of lenses, the eye, or of stimuli along the optic nerve. This proposal is difficult, but challenging. Physicists are trained in a mathematical framework, the outlook of physiologists is medical, while psychologists move largely in a humanistic environment. Moreover, these three groups of scientists differ in training and in their use of technical terms. Therefore, Ronchi implies that the present division of optics among the sciences not only hampers



**NEW and
completely revised!**

**14 Plates — Folding Map
56 Text Illustrations
Over 690 Pages**

Twenty-two world-famous specialists have participated to make this volume a classic. Editor Taylor has contributed six chapters and a glossary.

Partial List of Contents

- GEOGRAPHY IN THE NINETEENTH CENTURY
G. Tatham
- THE FRENCH SCHOOL OF GEOGRAPHY
R. J. Harrison Church
- THE GERMAN SCHOOL OF GEOGRAPHY
S. van Valkenburg
- THE WEST SLAV GEOGRAPHERS
J. Kral and J. Kondracki
- THE PROGRESS OF GEOMORPHOLOGY
S. W. Wooldridge
- GEOGRAPHICAL ASPECTS OF METEOROLOGY
F. K. Hare
- SOILS AND THEIR GEOGRAPHICAL SIGNIFICANCE
D. F. Putnam
- GEOGRAPHY AND ARCTIC LANDS
A. L. Washburn
- GEOGRAPHY AND REGIONALISM
E. W. Gilbert
- THE SOCIOLOGICAL ASPECTS OF GEOGRAPHY
J. W. Watson
- GEOGRAPHY AND AVIATION
Ellsworth Huntington
- GEOGRAPHY IN PRACTICE IN THE FEDERAL GOVERNMENT, WASHINGTON
John K. Rose
- GEOPOLITICS AND GEOPACIFICS
Griffith Taylor
- CARTOGRAPHY
W. W. Williams
- THE GEOGRAPHICAL INTERPRETATION OF AIR PHOTOGRAPHY
F. Walker
- THE RELATIONS OF GEOGRAPHY AND HISTORY
H. C. Darby

\$10

Order today from your favorite Bookseller or

PHILOSOPHICAL LIBRARY

Publishers

15 East 40 Street, Dept. A-54, New York 16, N. Y.
Expedite shipment by prepayment

Books on SATELLITES

Missiles Space Travel

- THE MAKING OF A MOON
by Arthur C. Clarke, 1957..... \$4.00
- REALITIES OF SPACE TRAVEL
edited by L. J. Carter, 1957..... \$7.00
- MAN INTO SPACE
by H. Oberth, 1957..... \$4.50
- ROCKETS, MISSILES, AND SPACE TRAVEL
by Willy Ley, 1957..... \$6.75
- GUIDED WEAPONS
by E. Burgess, 1957..... \$5.00
- ROCKET
by Sir Philip Joubert de la Ferte, 1957 \$3.00
- SCIENTIFIC USES OF EARTH SATELLITES
edited by J. A. Van Allen, 1956..... \$10.00
- ANNALS OF THE INTERNATIONAL GEO-PHYSICAL YEAR
Vol. I — Introductory Volume to IGY \$17.00
- ANNALS OF THE INTERNATIONAL GEO-PHYSICAL YEAR
Vol. II — Instruction Manual for Ionospheric Studies in the IGY.... \$17.00
- SATELLITE
by Erik Bergaust and William Beller, 1956 \$3.95
- THE INHABITED UNIVERSE
by K. W. Gatland, 1957..... \$3.75
- A TECHNICAL DICTIONARY OF ROCKETS AND ASTRONAUTICS
by G. Partel, 1955..... \$3.00

Write for information on our rocket and satellite slide sets.

Educational Division

**GENERAL ASTRONAUTICS
CORPORATION**

Dept. BS-3, 11 West 42nd St., New York 36, N. Y.

progress in understanding vision but propagates error. He gives many convincing examples of such errors.

Ronchi calls the end product of what our eyes perceive the *effigy*, not the image. He explains the distinction between them: "The image, whether real or virtual, defined as the locus of the centers of the waves emerging from an optical system, is a purely mathematical entity, entirely distinct from the figure seen. Images have a definite position; the figures seen are created and located by the observer, and may be placed by one observer in one way, by another observer in another way. An image is a mathematical entity, the figure seen is a psychological entity. To have identified these two en-

tities was a profound philosophical blunder. To convince millions of people that the two things are the same is one of the most ridiculous aspects of the teaching of science."

This contention is illustrated by a great many examples of effigies obtained with the aid of plane and spherical mirrors, and convergent and divergent lenses. As an instance of the failure of ordinary geometrical optics to account for appearances, Ronchi asks the reader to hold his hand at varying distances of about eight to 24 inches from his eye. Although it should vary in apparent size by a factor of three, the hand seems to undergo a much smaller change.

In presenting his thesis, Ronchi tends to be overzealous, seeming to portray the physicist's concept of optics as a villain. His contentions are, furthermore, not entirely new. The distinction between what we see and what we ought to see, according to the simple lens formulas, has been made by some philosophers, physicists, and teachers of science. In astronomy, "image" means different things to the visual observer and photographic astronomer. To the latter it is a particular distribution of silver grains on a plate, detectable impersonally by a microphotometer. We can distinguish a good photographic image from a poor one, and can judge the resolution by an optical system without once invoking the "effigy" concept which Ronchi holds paramount.

Because the science of vision, as considered in this book, runs directly into the difficult philosophical problems of consciousness, the author encounters semantic difficulties. There are many places where Ronchi, in his anxiety to prove a point, is not strictly fair to geometrical optics.

Despite minor lapses of this kind, wordiness, and some overstatement, this book is a refreshing and thought-provoking critique of the concepts of optics. Teachers will find it helpful, for it certainly clarifies many optical ideas often hard to convey in the classroom. Furthermore, it contains a fine historical survey of optics.

Edward Rosen, associate professor of history, College of the City of New York, ably translated the work from the original Italian.

J. ALLEN HYNEK
Smithsonian Astrophysical Observatory

Books on Astronomy

AMATEUR ASTRONOMER'S HANDBOOK, by J. B. Sidgwick.....	\$12.50
OBSERVATIONAL ASTRONOMY FOR AMATEURS, by J. B. Sidgwick.....	\$10.00
THE MOON, by Wilkins and Moore.....	\$12.00
THE SUN, by G. Abetti.....	\$12.00
THE AMATEUR ASTRONOMER, by P. Moore.....	\$4.50
APPLIED OPTICS AND OPTICAL DESIGN, by A. Conrady.....	\$2.95
Norton's STAR ATLAS.....	\$5.25
Webb's ATLAS OF THE STARS.....	\$6.50
SKALNATE PLESO ATLAS.....	\$5.25
BONNER DURCHMUSTERUNG.....	\$100.00
New: DISCOVERY OF THE UNIVERSE, by G. de Vaucouleurs.....	\$6.00

Write for free list. Out-of-print books located in a special search service. Books on telescope making and optical glass working. All books advertised and reviewed in Sky and Telescope.

HERBERT A. LUFT

NEW ADDRESS: 69-11 229th Street,
Oakland Gardens 64, New York

Sky and
TELESCOPE

Book Shelf

DISCOVERY OF THE UNIVERSE

by G. de Vaucouleurs \$6.00

a fascinating, strikingly illustrated history of the evolution of astronomical theory and methodology by the author of *Physics of the Planet Mars*.

The Macmillan Company

60 FIFTH AVENUE, NEW YORK 11, N.Y.



GIANT

Blow-ups of celestial phenomena for clubroom, den, school, and observatory. Galaxies, nebulae, moon enlarged to dramatic size, for first time at nominal cost. Sizes to 6' x 8'. Send for catalog.

2 x 3 foot blow-up of moon shipped in tube \$5.95
parcel post prepaid. No C.O.D.

ASTRO MURALS 231 W. 58th St., New York 19, N. Y.

TRANSACTIONS OF THE INTERNATIONAL ASTRONOMICAL UNION, Vol. IX, P. Th. Oosterhoff, editor, 1957. Cambridge University Press. 802 pages. \$15.00.

The ninth general assembly of the IAU, held at Dublin in 1955, has included only its main proceedings in this volume, with symposiums being published separately. This change in arrangement allows the *Transactions* more space for IAU commission reports, which give a very complete survey of progress in all branches of astronomy since the 1952 meeting in Rome.

Also in this volume are the minutes of the general sessions of the IAU at Dublin, all texts of the resolutions adopted, and the statutes and membership list of the union. The text is largely in English, with some sections in French.

NON-STABLE STARS, George H. Herbig, editor, 1957. Cambridge University Press. 200 pages. \$5.50.

The International Astronomical Union held several symposiums at its 1955 general assembly in Dublin, which are being reported in a series of books separate from the IAU *Transactions*. No. 3 of this series is entitled *Non-Stable Stars*, and is edited by a leader in the field, Dr. George H. Herbig, of Lick Observatory.

In this volume are published, in English, 23 contributions covering instability in low-luminosity stars of both late and early types, novae, hot stars of rather high luminosity, late-type variable stars, and close binary systems. Theoretical aspects of stellar instability are also discussed, and the book contains several Soviet contributions not circulated at Dublin.

For you and your friends . . .

make it an

ASTRONOMICAL CHRISTMAS

with these Sky Publications



Just Being Imported!

New Edition in Five Colors!

De Luxe ATLAS OF THE HEAVENS

by A. Becvar and his coworkers at the Skalnaté Pleso Observatory, Czechoslovakia

A striking advance in star atlases has been achieved in Antonín Becvar's new and improved ATLAS COELI 1950.0. It is handsomely printed in many colors: blue for the Milky Way, yellow for star clusters, red for galaxies; also green for planetaries and diffuse nebulae, with gray for dark nebulosities.

This use of color brings out dramatically such important astronomical facts as the concentration of star clusters along the Milky Way, the association of dark and bright nebulosity, and the rich cloud of galaxies in Coma Berenices and Virgo. Another new feature — over a hundred cosmic radio sources are plotted!

More than 35,000 celestial objects are shown on 16 charts covering the entire sky. All stars to as faint as magnitude 7.75 are included, and the data on variable and double stars have been brought up to date. Amateur and professional astronomers alike will welcome the addition of all the Bayer-letter and Flamsteed-number designations of the stars.

Now the ATLAS OF THE HEAVENS is clothbound, 16½ by 23 inches over-all, with the colored key on a handy flap. A transparent grid aids in reading star positions.

Order directly from Sky Publishing Corporation, distributors of the
Skalnaté Pleso Atlas of the Heavens for the Western Hemisphere.

De Luxe ATLAS OF THE HEAVENS..... \$9.75 postpaid

Regular edition (black and white only)..... \$5.25 postpaid

(The regular edition, with the same basic maps as the de luxe edition, is available at this reduced price, with transparent grid. Unbound and easily marked or written upon, the regular edition provides perfect worksheets for use at your desk or at the telescope.)

GIFT SPECIALS!

Photo Sets

1. **MOON PACKAGE:** For lunar enthusiasts, give Moon Sets and Lunar Crescent Sets, which together show a breathtaking view of the visible side of the moon. Moon Sets contain 18 large photographs made from unsurpassed Lick Observatory negatives of the first and last quarters. Lunar Crescent Sets are a matching series of 10 pictures from Lick Observatory, but for the waxing crescent 4½ days after new moon, and the waning crescent about five days before new moon. Each set is suitable for framing, or for use as an atlas. In this special offer, you will also receive the popular Lunar Map, which identifies a large number of features on the moon.

Moon Sets, Lunar Crescent Sets, and Lunar Map
A retail value of \$5.75 for \$5.00

2. **SKY PACKAGE:** With both amateur and professional astronomers, Sky Sets I and Sky Sets II have long been favorites. Each set contains 24 large photographs of astronomical wonders, including objects in the solar system and in the Milky Way, portraits of other galaxies, many made with the 200-inch telescope, and drawings of the Hale reflector by Russell W. Porter.

Sky Sets I and Sky Sets II
A retail value of \$8.00 for \$7.00

These gift specials on photo sets and books are good only
in the Western Hemisphere. Offers expire January 31, 1958.

Books

1. **BIG BOOKS:** This package of three popular books belongs on every amateur's book shelf. Each covers a different aspect: *Insight into Astronomy* is a general introduction for the layman and beginner; *Making Your Own Telescope* tells how to make and mount a 6-inch reflector at low cost; *The History of the Telescope* is the first book to tell the full story of the evolution of the telescope.

The History of the Telescope, by Henry C. King
Making Your Own Telescope, by Allyn J. Thompson
Insight into Astronomy, by Leo Mattersdorf

A retail value of \$20.00 for \$15.50

2. **SMALL BOOKS:** Here's an attractive package of our smaller publications. *Splendors of the Sky*, a 36-page picture book, is designed to help the beginner. The Lick Observatory *120-inch Album* records the complete story of the building of the world's second largest telescope. The *Lunar Map* identifies over 300 of the most important features on the moon. You also choose either of two informative booklets: *The Story of Cosmic Rays* or *Relativity and Its Astronomical Implications*.

Splendors of the Sky *120-inch Album*
Cosmic Rays or Relativity *Lunar Map*

A retail value of \$2.35 for \$2.00

A magazine on man's greatest adventure!

SPACEFLIGHT

Here is a popular, yet authoritative magazine on rockets, astronautics, and space-travel astronomy, written especially for the layman, and edited by members of the British Interplanetary Society. Leading authorities provide a comprehensive coverage of all the fields of science that play such an important part in this thrilling adventure of mankind. Rocketry, space medicine, atomic fuels, radar controls, the exploration of the planets are all treated in nontechnical language.

Spaceflight is printed during October, January, April, and July, volume 1 beginning with October, 1956. Be sure to specify the first issue for starting your subscription.

Subscription in United States and possessions, Canada, and Mexico: \$2.50, four issues (one year); \$4.50, eight issues (two years); \$6.00, twelve issues (three years). Single sample copy, 75 cents.

All items sent postpaid. Please enclose your check or money order payable to

SKY PUBLISHING CORPORATION

Harvard College Observatory

Cambridge 38, Massachusetts

And don't forget to give

SKY AND TELESCOPE

Here's an excellent gift for anyone on your Christmas list, from the budding young astronomer to the armchair scientist — a subscription to *Sky and Telescope*, the world's largest astronomical magazine, now printing 26,000 copies monthly. He will remember your thoughtfulness each month as he reads about this rapidly expanding science. A gift card, informing the recipient, will be sent in time for Christmas on orders received by December 10th. (Another way to give this magazine is through membership in an astronomical society that takes the group rate plan; consult your local club.)

Subscription in United States and possessions: \$5.00, one year; \$9.00, two years; \$13.00, three years. In Canada and Pan American Postal Union countries: \$6.00, one year; \$11.00, two years; \$16.00, three years. In all other countries: \$7.00, one year; \$13.00, two years; \$19.00, three years.



INFRARED SNIPERSCOPE TELESCOPE AND PARTS



See in the dark — without being observed. War-surplus Sniperscope M-2. Gov't. cost about \$1200. Used for industrial plant security, research lab experiments, infrared photography, spectroscopy, and the like. Instrument complete, ready to use. Includes power pack, infrared light source. Will operate from 6-volt auto battery. Battery or transformer available.

Stock #85,053-Y.....\$150.00 f.o.b.
(Shipping wt. approx. 12 lbs.) Barrington, N. J.

Save still more money! Build your own Sniperscope! We will furnish instructions and parts, including power packs, 1P25A image tubes, light units, filters, and the like. For details, request FREE Bulletin A-26-Y.

3" ASTRONOMICAL REFLECTOR

60- to 120-Power — An Unusual Buy!




Assembled — ready to use! See Saturn's rings, the planet Mars, huge craters on the moon, star clusters, moons of Jupiter, double stars, nebulae, and galaxies! Equatorial-type mounting with lock on both axes. Aluminum and over-coated 3" diameter f/10 primary mirror, ventilated cell. Telescope comes equipped with a 60X eyepiece and a mounted Barlow lens, giving you 60 to 120 power. A finder telescope, always so essential, included. Sturdy, hardwood, portable tripod.

Free with scope: Valuable STAR CHART and 272-page ASTRONOMY BOOK.

Stock #85,050-Y.....\$29.50 f.o.b.
(Shipping wt. 10 lbs.) Barrington, N. J.

4" REFRACTING TELESCOPE 240-POWER



Complete with Finder, Equatorial Mounting, Tripod, Eyepiece Extension, Star Diagonal, and Three Eyepieces

Our newest development. Built for rugged use, quality performance. Mounting made from heavy iron castings with machined bearings for smooth operation. Tripod has extra-heavy 60" hardwood legs. Telescope's weight is 42 lbs., giving stable, steady viewing. Big 4" objective is an air-spaced achromat, each element coated on both sides for low reflection. Three eyepieces supplied give you 48X, 120X, and 240X. Special Barlow lens also gives up to 500X. Star diagonal included for comfortable viewing at high angles. Rack-and-pinion focusing. All metal parts are plated to prevent rusting. Finder is 8 power. The usual price for a 4" refractor of comparable quality is over \$400, so our new model saves you almost 40%.

Stock #85,038-Y.....\$247.00 f.o.b.
(Shipping wt. 55 lbs.) Barrington, N. J.

SILICONE POLISHING AND CLEANING CLOTHS



Modern research produced this amazing new product. These cloths combine the miracle of silicones with a nonwoven, all-rayon cloth. Washable (use over and over) — they will dust, polish, and protect mirrors, reflectors, lenses, coated optics, crystal glass, plexiglass, lucite, and many other plastics. Vastly superior to diaper rags, cheesecloth, lens tissue. These are the only cloths that will not scratch any polished or smooth surface. They are lintless, nonoily, and nongreasy. Antifogging of glass is another benefit from their use. Cloths are 18" x 16".

Stock #60,059-Y.....Package of 3 cloths \$1.00 ppd.
Stock #70,137-Y.....Package of 100 cloths \$26.50 ppd.

Lower prices on larger quantities

ROTATING STAR CHART

Planisphere with a rotating chart shows well over 500 stars in the heavens and their relationships to each other at any selected day and hour. Table on reverse side supplies valuable information on constellations, navigation stars, locations of the planets according to month and year, dates of meteor showers, the zodiac, and the like.

Stock #9227-Y.....\$.50 ppd.

WAR-SURPLUS TELESCOPE EYEPIECE



Mounted Kellner Eyepiece, Type 3. 2 achromats, F.L. 28 mm., eye relief 22 mm. An extension added, O.D. 1 1/4", standard for most types of telescopes. Gov't. cost \$26.50.

Stock #5223-Y.....\$7.95 ppd.



EDSCORP SATELLITE TELESCOPE

OPTICS: The Satellite Scope has two important optical characteristics: A wide (51-mm.) diameter, low-reflection-coated objective lens. A six-element extremely wide-field, coated Erfle eyepiece that, in combination with the objective, gives 5.5 power with a big 12° field and over 7-mm. exit pupil.

OTHER USES FOR THE SATELLITE SCOPE

1. Makes a perfect wide-field finder. A special groove on the barrel helps in locating it in the finder mount. Fits our twin-ring finder mount, Stock No. 70,079-Y—\$9.95. 2. Use the Erfle eyepiece on your regular astronomical telescope. You will need our adapter, Stock No. 30,171-Y—\$3.95, which gives you an O.D. of 1 1/4". This eyepiece cost the government \$56.00! 3. Makes a wonderful comet seeker; see complete asterisms. 4. Makes a fine rich-field telescope; see wide areas of sky with deep penetration.

Especially Made for Members of MOONWATCH

Stock #70,074-Y.....\$49.50 ppd.

THE METZGER GLARE-REDUCTION SCREEN

The Metzger Glare-Reduction Screen is an accessory to refractors and reflectors, designed to cut planetary glare and help in observing finer planetary detail. This seemingly simple diffraction screen is a modern-type optical stop that sharpens the image without as much light loss as the old traditional circular stop. It has been most successful on the planet Mars, but is highly satisfactory on Jupiter, Saturn, Venus, and other planets. It was not designed especially for lunar observations because of the immense amount of light from the moon.

Much time and testing has been expended experimenting with finer and coarse screens to determine the most satisfactory. This mesh is the optimum. Obviously with the use of any diffraction screen, you will observe rays around any bright object in the field, but the object being observed will itself be in a darkened area. Since scattered light from a bright planet helps reduce contrast, the Glare-Reduction Screen improves contrast. The screen also helps when seeing conditions are not the best.

Before being placed on the market, the Glare-Reduction Screen was field-tested over a long period and is now being used by many amateur astronomers who have attested its value for critical observational work. The kraft-paper mount affords ample protection to the screen, which should be flat and should not be kinked, dented, or wavy. The screen does not have to fit snugly over the telescope tube but, if the fit is too loose, build up the mount with additional cardboard or corrugated packing. The Glare-Reduction Screen on 6" telescopes should center on the optical axis, which will align the cut-out hole.

Stock #70,138-Y—for 5" O.D. tubes.....\$2.95 ppd.
Stock #70,139-Y—for 7" O.D. tubes.....\$3.95 ppd.

ASTRONOMICAL TELESCOPE TUBING

Stock No.	I.D.	O.D.	Lgth.	Description	Price
80,038-Y	4 7/8"	5 1/4"	46"	Spiral-wound, paper	\$2.50
85,008-Y	6 7/8"	7 3/8"	60"		4.00
85,011-Y	2 7/8"	3"	48"		6.00
85,012-Y	3 7/8"	4"	60"	Aluminum	8.75
85,013-Y	4 7/8"	5"	48"		9.00
85,014-Y	6 7/8"	7"	60"		15.00

All tubing is shipped f.o.b. Barrington, N. J.

EDMUND SCIENTIFIC CO.

WAR-SURPLUS KELLNER EYEPiece

Mounted—Ready to Use—1 1/4" Outside
Diameter—Coated—1 1/4" Focal Length

Consists of an achromatic eye lens and an achromatic doublet field lens (Gov't. cost about \$30). The clear aperture of the lenses is approximately 1", giving wide exit pupil and a clear image. Excellent for any telescope when low power and a wide field are needed. Try it for 10 days—if you don't agree that the performance is better than any commercial type selling for two or three times our price, we will refund your money in full.



Stock #50,130-Y.....\$5.90 ppd.

Rack & Pinion Eyepiece Mounts



For Reflectors



For Refractors

Now you can improve performance in a most important part of your telescope—the eyepiece holder. Smooth, trouble-free focusing will help you to get professional performance. Look at all these fine features: real rack-and-pinion focusing with variable tension adjustment; tube accommodates standard 1 1/4" eyepieces and accessory equipment; lightweight aluminum body casting; focusing tube and rack of chrome-plated brass; body finished in black wrinkle paint. No. 50,077-Y is for reflecting telescopes, has focus travel of over 2", and is made to fit any diameter or type tubing by attaching through small holes in the base. Nos. 50,103-Y and 50,108-Y are for refractors and have focus travel of over 4". Will fit our 2 7/8" I.D. and our 3 7/8" I.D. aluminum tubes respectively.

Stock #50,077-Y (less diagonal holder) \$9.95 ppd.
Stock #60,035-Y (diagonal holder only) 1.00 ppd.
Stock #50,103-Y (for 2 7/8" I.D. tubing) 12.95 ppd.
Stock #50,108-Y (for 3 7/8" I.D. tubing) 13.95 ppd.

HUYGENS EYEPieces

Here are some really terrific values in eyepieces! The three eyepieces listed below are manufactured by one of the world's best producers of optical components. We have searched the world's markets, including Germany and France, to find a real quality eyepiece. The image clarity, the workmanship evidenced in the metal parts, will prove the skill and experience of Goto Optical Company, Tokyo. Guaranteed terrific buys!

HUYGENS TYPE—STANDARD 1 1/4" DIAMETER
6-mm. (1/4") Focal Length
Stock #30,063-Y.....\$8.50 ppd.


12.5-mm. (1/2") Focal Length
Stock #30,064-Y.....\$8.00 ppd.

COMBINATION EYEPiece—10-mm. and 20-mm.
Stock #30,065-Y.....\$9.00 ppd.

7X FINDER TELESCOPE—ACHROMATIC

Stock #50,080-Y Finder alone, less ring mounts \$9.95
Stock #50,075-Y Ring mounts per pr.\$3.95


RUBBER PITCH-LAP MAT Saves Mirror Makers Time and Trouble




A rubber pitch-lap mat is used for forming the pitch-lap channels. Just pour the melted pitch on the tool, lay the mat on top and then press it in with your mirror to form the channels. When the pitch sets, an easy pull will remove the mat. In minutes instead of hours you are ready to polish. Eliminates time-consuming and tedious hand-cutting of the channels of the pitch lap. All those disappointing break outs of the pitch, common when hand-cutting the channels, are avoided. You no longer have to pour and cut two or three laps before getting a usable one. With our Rubber Pitch-Lap Mat you can use the first one you make.

For
Stock No. Size Mirror Dia. Price ppd.
50,171-Y 13 1/2" x 13 1/2" 8", 10", 12" \$2.00
60,061-Y 6" x 6" 4 1/4", 6" 1.00

DOUBLE AND TRIPLE YOUR TELESCOPE'S POWER WITH A BARLOW LENS



WHAT IS A BARLOW? A Barlow lens is a negative lens used to increase the power of a telescope without resorting to short focal length eyepieces, and without the need for long, cumbersome telescope tubes. Referring to the diagram above, a Barlow is placed the distance P inside the primary focus of the mirror or objective. The Barlow diverges the beam to a distance Q. This focus is observed with the eyepiece in the usual manner. Thus, a Barlow may be mounted in the same tube that holds the eyepiece, making it very easy to achieve the extra power. The new power of the telescope is not, as you might suppose, due to the extra focal length given the objective by the difference between P and Q. It is defined as the original power of the telescope times the quotient of P divided into Q!



Beautiful chrome mount. We now have our Barlow lens mounted in chrome-plated brass tubing with special spacer rings that enable you to vary easily the power by sliding split rings out one end and placing them in other end. Comes to you ready to use. Just slide our mounted lens into your 1 1/4" I.D. tubing, then slide your 1 1/4" O.D. eyepieces into our chrome-plated tubing. Two pieces provided, one for regular focal length eyepieces and one for short focal length ones.

Remember, in addition to doubling and tripling your power, a Barlow lens increases your eye relief and makes using a short focal length eyepiece easier.

Don't fail to try one of these. Many people do not realize the many advantages of a Barlow and the much greater use they can get from their telescopes. Our Barlow has a focal length of -1-5/16". We have received many complimentary letters about this lens. So sure are we that you will like it that we sell it under a 30-day guarantee of satisfaction or your full purchase price returned—no questions asked. You can't lose, so order today.

Stock #30,200-Y Mounted Barlow lens\$8.00 ppd.

ATTENTION—BEGINNERS AND JUNIOR ASTRONOMY CLUBS

To solve the problem of low-cost lenses of good quality for a beginner's telescope we have had these achromatic lenses designed and mass produced. They are swell!

ACHROMATIC OBJECTIVE, 32-mm. diameter (1 1/4"), F. L. 31". (Will get 31X with 1" eyepiece; 62X with 1/2" eyepiece.)

Stock #30,201-Y.....\$2.50 ppd

ACHROMATIC OBJECTIVE, 42-mm. diameter (1-21/32"), F. L. 41". (Will get 41X with 1" eyepiece; 82X with 1/2".)

Stock #30,202-Y.....\$4.00 ppd.

PRISM STAR DIAGONAL

For comfortable viewing of the stars near the zenith or high overhead with refracting telescopes using standard size (1 1/4" O.D.) eyepieces, or you can make an adapter for substandard refractors. Contains an excellent quality aluminized right-angle prism. Tubes are satin chrome-plated brass. Body is black wrinkle cast aluminum. Optical path of the system is about 3 1/2".



Stock #70,077-Y.....\$12.00 ppd.

SPITZ MOONSCOPE

A precision-made 32-power reflecting telescope—by makers of Spitz Jr. Planetarium. Clearly reveals the craters of the moon, shows moons of Jupiter, other wonders of the heavens. Based on same principles as world's giant telescopes. Stands 56" high on removable legs. Adjustable 3" polished and corrected mirror. Fork-type altazimuth mount rotates on full 360° circle—swings to any location in the sky. Fascinating 18-page instruction book; sturdy carrying case.



Stock #70,068-Y.....\$14.95 ppd.

"MAKE-YOUR-OWN" 4 1/4" MIRROR KIT

The same fine mirror as used in our telescopes, polished and aluminized, lenses for eyepieces, and diagonal. No metal parts.

Stock #50,074-Y.....\$16.25 ppd.

6X FINDER TELESCOPE



Has crosshairs for exact locating. You focus by sliding objective mount in and out. Base fits any diameter tube—an important advantage. Has 3 centering screws for aligning with main telescope. 20-mm. diam. objective. Weighs less than 1/2 pound.

Stock #50,121-Y.....\$8.00 ppd.

MISCELLANEOUS ITEMS

KELLNER EYEPiece—2" focal length (1 1/4" O.D.). Mount of black anodized aluminum.

Stock #30,189-Y.....\$6.00 ppd.

60° SPECTROMETER PRISM—Polished surfaces 18-mm. x 30-mm.—flat to 1/2 wave length.

Stock #30,143-Y.....\$8.25 ppd.

BRASS TUBING

2 pieces, 3" long, slide fitting. Blackened brass. I.D. 1-3/16", O.D. 1-5/16".

Stock #40,165-Y.....\$1.75 ppd.

30-day Money Back Guarantee
as with all our Merchandise!

BE SURE TO GET FREE CATALOG "Y"

Fantastic variety—never before have so many lenses, prisms, optical instruments, and components been offered from one source. Positively the greatest assembly of bargains in all America. Imported! War Surplus! Hundreds of other hard-to-get optical items. Write for Free Catalog "Y."

ORDER BY STOCK NUMBER . . . SEND CHECK OR MONEY ORDER . . . SATISFACTION GUARANTEED!

BARRINGTON • NEW JERSEY

PANGRO mirrors, inc.
PANGRO mirrors, inc.

Reg. U.S. Pat. Off.

Your mirror may now have the same aluminum and quartz over-coating, with high reflection, durability, and guaranteed permanent adhesion, that we are giving Cave Optical Co. for the Astrola telescopes. Your mirror will be carefully packed and on the way back to you within 24 hours.

6-inch.....	\$6.50	10-inch.....	\$11.25
8-inch.....	\$8.50	12-inch.....	\$16.00

Prices are f.o.b. Los Angeles.

Since 1933

PANGRO MIRRORS, INC.
 Research and Production Laboratories
 2958 Los Feliz Blvd., Los Angeles 39, Calif.

LOW-EXPANSION ELLIPTICAL DIAGONALS
 Aluminized, 1/8 wave.

Minor Axis		
1/2"	3.75	
3/4"	3.75	
1 1/4"	5.00	
1 1/2"	6.50	

MODIFIED RAMSDEN EYEPIECES
 1/2", 3/4", 1 1/4", 2", 3", and 4" focal lengths.
 Attractively mounted in 1 1/4" O.D. silicon-aluminum bushings.

\$5.50 each, \$14.00 per set (any 3)

VERY FINEST WASHED ROUGE for the final stages of polishing and figuring.
 In 4 oz. jar (wet).....\$1.80

Write for free catalog.

NYE OPTICAL CO.
 2100 Cherry Ave., Long Beach 6, Calif.

TELESCOPE MAKERS
Everything for the Amateur KITS.....\$4.50 up

Thousands of our customers with no special training have built powerful precision telescopes with our quality supplies, instructions, and guidance.

Send for Complete Instructions, 10c

ALUMINIZING
 Superior Reflecting Surface.
 Guaranteed not to peel or blister.

MIRRORS MADE TO ORDER
MIRRORS TESTED FREE

EYEPIECES CELLS | **PRISMS BINOCULARS**

PRECISION TRI-TURRET

Holds 3 standard 1 1/4" O.D. eyepieces. Smooth turn to grooved notch aligns eyepiece precisely, ready to focus for various powers. Suitable for reflectors or refractors. **\$15.75**




ORTHOSCOPIC EYEPIECES
HIGHEST QUALITY • MADE IN U.S.A.
 Special four-element design, with fluoride-coated lenses, gives a wide flat corrected field. Standard 1 1/4" O.D. — E.F.L. 6-8-12-16-24-mm. Postpaid **\$15.95**

Write today for FREE Catalog.

Precision Optical Supply Co.
 1001-H East 163rd St., New York 59, N. Y.

GLEANINGS FOR ATM's

CONDUCTED BY ROBERT E. COX

A NEW TEST FOR CASSEGRAINIAN SECONDARIES

OPTICAL THEORY teaches that a spherical wave front converges to a point focus and forms an image free from longitudinal aberration. It also makes clear that this condition cannot be obtained with any single spherical surface receiving parallel light. It can, however, be obtained with an aspherical refracting surface.

The required shape of such a surface depends upon its radius and the indexes of refraction of the two media that it separates, n for the first medium, n' for the second:

A. If light passes from a denser to a less dense medium (as glass to air), the surface must be hyperboloidal, and its eccentricity will be n/n' .

B. If light passes from a less dense to a denser medium (air to glass), the surface must be ellipsoidal, with an eccentricity of n/n' .

These two situations are illustrated in Fig. 1 for both concave and convex surfaces, where the media are air and BSC-2 glass with an index of 1.517. The statements above can be verified by trigonometric ray tracing.

Although I came upon these ideas inde-




Fig. 2. A sketch of the optics of a Cassegrainian reflecting telescope.

pendently in the course of my hobby of ray tracing and telescope making, I later learned from Dr. James G. Baker that they date back to the 17th-century French mathematician, René Descartes, and his investigation of the curves known as Cartesian ovals. These ideas are well explained in M. von Rohr's book, *Theory of the Formation of Images in Optical Instruments*, pages 24 to 26 of the English translation of 1920.

What practical use can be made of this principle? One obvious application is a solution to the difficult problem of making and testing a hyperboloidal secondary mirror for a Cassegrainian telescope. I shall describe here how one can make and test a plano-convex lens, to be figured in the test stages, and the convex surface of this same lens, when aluminized, serves as the secondary mirror!

In Fig. 2 is sketched the Cassegrainian




Fig. 1. Various cases of a principle of optics applied by Burt A. Norman in testing a Cassegrainian secondary.




Fig. 3. The test setup as used when an optical flat is available.

optical system, and in Fig. 3 the testing arrangement. The secondary is tested from the convex side by autocollimation. If a good optical flat is not available for this test, simply silver the plane surface of the lens. Even if the back is not used as a reflecting surface, it should be flat to less than 1/8 wave; polish need not be perfect.

For a Cassegrainian whose secondary mirror is located the distance P inside the primary focus, the secondary's back focal length being P' , the following relationships are well known:

Radius of curvature of central zone

$$R = 2PP'/(P' - P),$$

Eccentricity of hyperboloid

$$e = (P' + P)/(P' - P).$$


The design is chosen so that the eccentricity is numerically equal to the refractive index of some easily obtainable type of glass. From this glass make a plano-convex lens having a radius of curvature equal to R , and a paraxial back focal

Instead of the usual off-axis Foucault setup, use a beam splitter to insure that the light source and knife edge lie exactly on the optical axis of the lens. Figure the convex surface of the lens (changing it from a sphere) until a null test is obtained. This means that the knife-edge cutoff produces an even darkening all over the surface at one time, just as if a spherical mirror were being tested. But with this test, the refractive index of the


There is a practical restriction on the application of this testing method to a particular telescope. The factor by which the secondary mirror increases the focal length of the primary is called the amplifying ratio, P'/P . The glass selected for the lens must match the design, having the proper refractive index for the wave length of the test light. For example, if sodium light is used for the test source, then a secondary made from BSC-2 glass must have an amplifying ratio of 4.87.

To find the paraxial focal length of the test lens from Fig. 4, read off the amplifying ratio and multiply this figure by **P**. For greater accuracy, use the formula

Catalogue No. 4 of the Hayward Glass Corp., 217 Magnolia Ave., Whittier, Calif.,



Right: Fig. 5. This chart from Eastman Kodak Co. gives the indexes of refraction for some common types of glass. The abscissa of the chart is the nu-value, or reciprocal dispersion; it is not involved in the Norman test, where monochromatic light must be used.



AMERICAN MADE

STANDARD MODEL "A",	6-inch	\$295.00
STANDARD MODEL "B",	8-inch	\$375.00
STANDARD MODEL "C",	10-inch	\$475.00

These instruments are fully portable—as they can be assembled or taken down in three minutes. Each comes with three of the finest oculars. The equatorial head and stand are of cast aluminum. The fiberglass tube is made by W. R. Parks. Optics are corrected to $\frac{1}{8}$ wave or better and are quartz coated. ASTROLAS will resolve double stars to the Dawes limit. Clock drives, rotating tubes, setting circles furnished at additional cost.

ARE YOU SATISFIED WITH YOUR PRESENT MIRROR?

If your mirror does not give the fine performance you need for satisfactory observing, the expert opticians at Cave Optical Co. will refigure it at a reasonable price, or offer you a new precision-ground pyrex mirror. Whichever you choose, your mirror will have a fine figure, good to $\frac{1}{8}$ wave length or better.

REFIGURING MIRRORS

Refiguring, aluminizing, and quartz coating your mirror. Diagonal also provided.

6-inch	\$35.00
8-inch	\$50.00
10-inch	\$75.00
12½-inch	\$120.00

PYREX MIRRORS


New pyrex mirrors, aluminized and coated. Diagonal also provided.

6-inch	\$60.00
8-inch	\$92.50
10-inch	\$160.00
12½-inch	\$250.00

*All prices f.o.b. our plant, Long Beach, Calif., and subject to change without notice.
Send for catalogue.*

CAVE OPTICAL COMPANY

4137 E. Anaheim St., Long Beach 4, Calif.



Standard Model "A" 6-inch ASTROLA, f/8,
complete with 3 oculars (72x, 180x, 315x)
\$295.00

*****DELUXE PYREX*****

Reflecting Telescope Kits

Our kits have PYREX mirror blank, PYREX tool the same thickness, ample supply of optical quality abrasives, fast polishing cerium oxide, red rouge and pitch. Packed in metal cans.

Size	Thickness	Price
4 1/4"	3/4"	\$ 6.00
6"	1"	\$10.50
8"	1 1/2"	\$18.75
10"	1 3/4"	\$33.65
12 1/2"	2 1/8"	\$59.95

ADD POSTAGE: 1st and 2nd postal zones from Detroit, add 5%; 3rd and 4th, add 10%; 5th and 6th, add 15%; 7th and 8th, add 20%. Or we will ship C.O.D.

Send for free catalog of supplies, accessories, and refracting telescopes.

ASHDOWNE BROS.

18450 Grand River Detroit 23, Michigan

Precision Diagonals

You will get the best possible performance from your telescope with one of our clear fused quartz diagonals. Accuracy guaranteed 1/20 wave.

Ellipse 1.25" x 1.77"	\$11.00
Ellipse 1.5" x 2.12"	\$14.00

Pyrex diagonals, 1/8 wave accuracy.

Ellipse 1.25" x 1.77"	\$ 5.00
Ellipse 1.5" x 2.12"	\$ 8.00

Aluminum coating \$1.00 extra.

Send for our complete list of supplies, quartz mirrors, blanks, oculars, coatings, and accessories.

E & W OPTICAL CO.

2406 E. Hennepin Ave.
Minneapolis 13, Minn.

lists various types of optical glass, with their indexes for different colors of light, such as the mercury green line and the sodium yellow line. Bausch and Lomb Optical Co., Rochester 2, N. Y., also makes optical glass. To be very exact, specify when ordering that you wish to know these indexes for the particular melt of glass, because each batch of glass is slightly different.

Choice of the light source for this test is of great importance. The old trick of rubbing salt into the wick of an oil lamp to get sodium light does not give enough brightness. A sufficiently bright mercury light source that costs less than five dollars, the General Electric F4T 5/BL black-light lamp, has been suggested to me by Robert E. Cox. Do not get the BLB lamp, for it has a coating that passes only invisible light and is useless as a test source.




Fig. 6. The arrangement of parts in an inexpensive mercury light source.




Fig. 7. The mercury light source disassembled. The black-light tube fits into a metal container seven inches long. At the front are the transformer and four-wire switch. Photo by Robert E. Cox.

The ballast is a General Electric 89G435 transformer. In the wiring diagram for the light source, Fig. 6, the four-wire on-off switch is an FLS 40-watt, 125-volt a.c. switch, obtainable from Sears Roebuck or other stores handling fluorescent lamps. The parts are seen in Fig. 7.

To isolate the green mercury line at 5461 angstroms, an Eastman Kodak Wratten 57A filter is recommended; a two-inch square of this material in gelatin costs 60 cents. The same filter can be obtained glass-mounted (grade B is satisfactory), if this more permanent form is worth the higher price.

Other monochromatic light sources that should be investigated are the Osram spectral lamps available from Edmund Scientific Co., Barrington, N. J. The helium, mercury, and sodium Osram lamps should be bright enough for testing the secondary mirror. They are listed at \$29.50, but other electrical components needed to energize these lamps will run the total cost to over \$45.

This test method is only one of the possible applications of the principle of Cartesian ovals to optical design, a principle which is less familiar to amateur telescope makers than it should be. Probably $e = n/n'$ can benefit the amateur optical worker in other ways.

BURT A. NORMAN
Franklin Center
Quebec, Canada

Read This Advertisement

Here is a combination of a Barlow and a particular ocular which gives outstanding results. It consists of our new Barlow and our 16.3-mm. (2/3" focal length) Erfle eyepiece. While the Barlow was not specifically designed to work with this eyepiece, it does so to an astonishing degree. All images are sharp and hard to the very edge of the field.

The Barlow gives magnification up to slightly over three times that of the ocular alone. It is achromatic, coated, and mounted to the U. S. standard size of 1.250 inches.

The modified Erfle eyepiece has a field of 75 degrees with excellent eye relief. The combination gives the equivalent focal length of slightly under 6 mm. Many users state it is far superior to any shorter focal length ocular of equivalent magnification.

The Barlow sells for \$16.00 postpaid, and the Erfle for \$14.75 postpaid. Both are guaranteed to perform as stated above or money refunded.

ORTHOSCOPIC OCULARS—All hard coated, standard 1 1/4-inch outside diameter.

28-mm.	\$15.00	10.5-mm.	\$16.25	4-mm.	\$17.25
16.3-mm. (Erfle) ...	\$14.75	7-mm.	\$17.25	Barlow 3x	\$16.00

Telescopes

1000 North Seward Street
Los Angeles 38, Calif.

Warranted to equal or surpass any oculars obtainable anywhere or money refunded.

Finished mirrors, mirror kits, spiders, elliptical flats, focusing devices, aluminizing.

Send for catalogue.

Magnusson Heavy-Duty Mounting

- Latitude adjustment.
- 1 1/4" x 17" declination shaft.
- Ball-bearing polar-axis shaft, 1 1/2" on north end —1" on south end x 12" long. Suitable for clock drive.
- High tensile aluminum black-finish castings, with solid-steel shafts.
- 11-pound cast-iron balance weight.
- Locking screws on polar and declination axles.
- 5" brass setting circles with pointers.
- Weight approximately 38 pounds complete.
- Custom-made saddle to fit your telescope.

State size of tube, up to 12". Tripod not included.
With setting circles \$125.00 f.o.b. Arvada
Without circles \$96.00 f.o.b. Arvada


SETTING CIRCLES

	Aluminum	Brass
5" circles, 1" holes, set of two	\$15.00	\$19.50
6" circles, 1 1/2" holes, set of two	\$17.00	\$22.10
8" circles, 1 1/2" holes, set of two	\$24.00	\$31.20

Add \$2.00 per set for change of hole sizes.

O. MAGNUSSON

14570 West 52nd Ave., Arvada, Colorado



ED. NOTE: The test proposed by Mr. Norman is not a familiar one to optical designers, and it may be appropriately called "the Norman test." Mr. Norman is a native of Arkansas, and has been a construction engineer at hydroelectric, chemical, and other industrial plants, as well as the Chalk River project in Ontario.

The well-known optical designer, Dr. James G. Baker, comments, "A fine practical application of a very old principle. I do not recall having seen this optical theorem used for a Cassegrainian test. This is a good example of the contribution that amateurs can often make to optics."

AIRSPACED OBJECTIVES

MOUNTED IN ALUMINUM CELLS f/15

We offer the lowest priced, hand-corrected, precision, American-made astronomical objective, mounted in a black anodized aluminum cell. Our reputation has been established over the years as the most reliable source of high quality astronomical lenses.

"Those in the know" BUY FROM US BECAUSE:

Each lens is thoroughly tested by us and is guaranteed to resolve two seconds of arc or better. They are corrected for the C and F lines (secondary chromatic aberration). The zonal spherical aberration and the chromatic variation of spherical aberration are negligible. The cell is machined to close tolerances so that it will fit directly over our standard aluminum tubing, eliminating any mounting problems.

3 1/4" diam., 48" f.l. (uncoated).....\$28.00 4 1/8" diam., 62" f.l. (uncoated).....\$60.00
Same as above with coating.....\$32.00 Same as above with coating.....\$69.00

We can supply ALUMINUM TUBING for the above lenses.

"BIG" ACHROMATIC TELESCOPE OBJECTIVES

We have the largest selection of diameters and focal lengths in the United States available for immediate delivery. These are perfect magnesium-fluoride coated and cemented Gov't. surplus lenses made of finest crown and flint optical glass. Not mounted. Fully corrected. Tremendous resolving power. They can readily be used with eyepieces of only 1/4" focal length, thereby producing high powers. Guaranteed well suited for astronomical telescopes, spotting scopes, and other instruments. Gov't. cost up to \$100.

Diameter	Focal Length	Each	Diameter	Focal Length	Each
54 mm (2 1/8")	254 mm (10")	\$12.50	83 mm (3 1/4")	660 mm (26")	\$28.00
54 mm (2 1/8")	300 mm (11.8")	12.50	83 mm (3 1/4")	711 mm (28")	28.00
54 mm (2 1/8")	330 mm (13")	12.50	83 mm (3 1/4")	762 mm (30")	28.00
54 mm (2 1/8")	390 mm (15.4")	9.75	83 mm (3 1/4")	876 mm (34 1/2")	28.00
54 mm (2 1/8")	508 mm (20")	12.50	83 mm (3 1/4")	1016 mm (40")	30.00
54 mm (2 1/8")	600 mm (23 1/2")	12.50	102 mm (4")	876 mm (34 1/2")	60.00
54 mm (2 1/8")	762 mm (30")	12.50	108 mm (4 1/4")	914 mm (36")	60.00
54 mm (2 1/8")	1016 mm (40")	12.50	110 mm (4 3/8") *	1069 mm (42-1/16")	60.00
54 mm (2 1/8")	1270 mm (50")	12.50	110 mm (4 3/8")	1069 mm (42-1/16")	67.00
78 mm (3-1/16")	381 mm (15")	21.00	128 mm (5-1/16") *	628 mm (24 3/4")	75.00
80 mm (3 1/8")	495 mm (19 1/2")	28.00	128 mm (5-1/16")	628 mm (24 3/4")	85.00
81 mm (3-3/16")	622 mm (24 1/2")	22.50			

● We can supply ALUMINUM TUBING for the above lenses. ●

COATED BINOCULARS



American Type



"Zeiss" Type

Beautiful imported binoculars, precision made, at a low, low price. Above we have pictured the two most popular types. The American Type offers a superior one-piece frame and a clean design, pleasing to the eye. Complete with carrying case and straps. Price plus 10% Federal tax.

SIZE	TYPE	C. FOCUS	IND. FOCUS
6 x 15	OPERA	—	\$12.75
6 x 30	"ZEISS"	\$18.75	16.75
7 x 35	"ZEISS"	21.25	19.25
7 x 35	AMERICAN	23.50	—
7 x 35	AMERICAN WIDE	—	—
	ANGLE 10°	37.50	—
7 x 50	"ZEISS"	24.95	22.50
7 x 50	AMERICAN	32.50	—
8 x 30	"ZEISS"	21.00	18.25
10 x 50	"ZEISS"	30.75	28.50
20 x 50	"ZEISS"	41.50	39.50

MONOCULARS




Brand new, coated optics, complete with pigskin case and neck straps.

	Price		Price
6 x 30	\$10.00	7 x 50	\$15.00
8 x 30	11.25	16 x 50	17.50
7 x 35	12.50	20 x 50	20.00

"MILLIONS" of Lenses, etc.
Free Catalogue

We pay the POSTAGE — C.O.D.'s you pay postage. Satisfaction guaranteed or money refunded if merchandise returned within 30 days.

"GIANT" 3" TELESCOPE



40 power

Special Price \$57.50

Never before has anything like this been offered at so low a price. Here is another example of American ingenuity. Big 3" diameter achromatic coated objective which will give needle-sharp crystal-clear images. Focusing is a delight with the micrometer spiral focusing drawtube. Light-weight aluminum construction throughout, black crackle finish, length open 22 inches, closed 17 inches. This telescope gives an upright image—it is WONDERFUL for astronomy, SUPERB for long distances, EXCELLENT as a spotting scope.



"GIANT" EYEPIECE

WIDE ANGLE ERFLE (68° Field) EYEPIECE. Brand new coated 1 1/4" E.F.L. Focusing mount. 3 perfect achromats, 1-13/16" aperture \$18.50

WIDE ANGLE ERFLE 1 1/2" E.F.L. Brand new; contains Eastman Kodak's rare-earth glasses; aperture 2"; focusing mounts; 65° field \$18.50

1 1/4" Diam. Adapter for above eyepieces \$9.95

LENS CLEANING TISSUE — Here is a wonderful Gov't. surplus buy of Lens Paper which was made to the highest Gov't. standards and specifications. 500 sheets size 7 1/2" x 11" \$1.00



MOUNTED EYEPIECES

The buy of a lifetime at a great saving. Perfect war-surplus lenses set in black anodized standard aluminum 1 1/4" O.D. mounts.

F.L.	TYPE	PRICE
12.5 mm (1/2")	Symmetrical	\$ 6.00
16 mm (5/8")	Erfle (wide angle)	12.50
16 mm (5/8")	Triplet	12.50
18 mm (3/4")	Symmetrical	6.00
22 mm (27/32")	Kellner	6.00
32 mm (1 1/4")	Orthoscopic	12.50
35 mm (1 3/8")	Symmetrical	8.00
55 mm (2-3/16")	Kellner	6.00
56 mm (2 1/4")	Symmetrical	6.00

COATED 75 cents extra.

ASTRONOMICAL MIRRORS

These mirrors are of the highest quality, polished to 1/4-wave accuracy. They are aluminized, and have a silicon monoxide protective coating. You will be pleased with their performance.

	Diam.	F.L.	Postpaid
Plate Glass	3-3/16"	42"	\$ 9.75
Pyrex	4 1/4"	45"	13.50
Pyrex	6"	60"	25.00

MIRROR MOUNT

Cast aluminum. Holds all our mirrors firmly with metal clips. Completely adjustable. Assembled, ready to use.

3-3/16" Mount fits our 4 1/2" tubing \$4.00 ppd.
4 1/4" Mount fits our 5" tubing 4.00 ppd.
6" Mount fits our 7" tubing 7.00 ppd.

Aluminum Telescope Tubing

O.D.	I.D.	Price Per Ft.
2 1/4"	2 1/8"	\$1.20 ppd.
3 3/8"	3 1/4"	1.75 ppd.
4 1/2"	4 3/8"	2.75 ppd.
5"	4 7/8"	2.75 ppd.
7"	6 7/8"	3.00 f.o.b.

Focusing Eyepiece Mounts Rack & Pinion Type

The aluminum body casting is finished in black crackle paint and is machined to fit all our aluminum tubing. Has a chrome-plated brass focusing tube, which accommodates standard 1/4" eyepieces.

For 2 1/8" I.D. Tubing Postpaid \$12.95
For 3 1/4" I.D. Tubing " 12.95
For 4 3/8" I.D. Tubing " 12.95


REFLECTOR TYPE FOR ALL SIZE TUBING:

Complete with diagonal holder \$ 9.95

Aluminum Lens Cells Black Anodized

Cell for Lenses	Cell Fits Tubing	Price
54 mm Diam.	2 1/8" I.D.	\$ 3.50
78 mm "	3 1/4" "	6.50
81 mm "	3 1/4" "	6.50
83 mm "	3 1/4" "	6.50
110 mm "	4 3/8" "	10.50

3X TELESCOPE



Makes a nice low-priced finder. Brand new; has 1" Achromatic Objective. Amici Prism Erecting System, 1 3/4" Achromatic Eye and Field Lens. Small, compact, wt. 2 lbs.

Gov't. cost \$200. \$9.75

FIRST SURFACE MIRRORS

Size	Postpaid	Size	Postpaid
14" x 16"	\$10.00	5 1/4" x 7 1/4"	\$3.00
10" x 10"	5.00	5" x 5"	2.00
9" x 11-3/16"	5.00	4" x 5"	1.85
8" x 10"	4.25	4" x 4"	1.50

All mirrors are 1/4" thick.

TELEVISION PROJECTION LENS

Brand New, f/1.9, E.F.L. 5 inches. Manufactured by Bausch & Lomb. We purchased entire lot of these discontinued units. Five elements, smallest lens 2", largest 4 1/8". Completely assembled 6" in length. All surfaces hard coated. Get this BARGAIN now.

ONLY \$22.50

A. JAEGER'S

• THE GLASS HOUSE •

691 S MERRICK RD. LYNBROOK, N.Y.



Japan's Finest Telescopes now available to World Astronomers



This is the 4" Equatorial Telescope which was recently manufactured and delivered to Mr. A.T. Graves of 9 Synswyck Road, Darien, Connecticut, famous industrialist and amateur astronomer, who placed a special order for its manufacture during his stay in Tokyo. That the quality of this telescope has come to be recognized can be seen from the recent increase in the numbers exported. For example, they were directly shipped to:

Sainte-Marie College
1180 Rue Bleury, Montreal 2, Canada.
Dr. S. von Numers
The Legation of Finland,
Ottawa, Canada.

Aside from direct exports many are now being shipped indirectly overseas through various sales agents to the United States and Latin America attesting to their growing world-wide popularity.

Established in 1926, Japan's
Oldest and Largest Firm
Specializing in Astronomical
Telescopes

Achromatic Objectives

For use in astronomical telescopes. Mounted in cell.

3" C.A. 47.2" F.L. \$61.50

4" C.A. 59.1" F.L. \$137.00

Oculars—Fit American size barrel.

Orthoscopic (regular type): 4, 6, 9, 12.5, 18, 25 mm. \$14.40 each

Kellner: 6, 9, 12.5, 18, 25, 40 mm. \$12.90 each

Huygens: 6, 9, 12.5, 18, 25, 40 mm. \$6.70 each

Star Diagonals

For use with eyepieces under 25 mm. \$10.90

Erecting Prisms

For use with eyepieces under 25 mm. \$11.90

Sun Projecting Screen Set \$11.90

(Prices including shipping costs)
The above prices also include the anticipated import duties so in making remittances 25% should be deducted from the listed prices.

GOTO OPTICAL MFG. CO.

1-115 SHIMMACHI, SETAGAYA-KU, TOKYO, JAPAN

Cable Address:

GOTOPTIC TOKYO

A VERSATILE OBSERVING SETUP WITH AN 8-INCH REFLECTOR

AFTER several years of making and observing with many small instruments, I have constructed an 8-inch reflector which combines fine optics with versatility and rigidity.

The heart of this telescope, an 8-inch $f/7.5$ paraboloidal mirror, is mounted in a hardwood cell similar to that described in Allyn J. Thompson's *Making Your Own Telescope*. The cell holds the mirror without flexure and has never needed realigning. The mirror is housed in a welded, cork-lined, 10" aluminum tube.

At first I used a circular spider to hold my elliptical flat, but since this was not perfectly rigid, I made a three-legged spider and used an antidiffraction mask on the mirror (*Sky and Telescope*, April, 1957, page 289). The arrangement gives fine refractor-like star images and insures alignment.


The equatorial mounting of $2\frac{1}{2}$ " pipe fittings has $1\frac{1}{2}$ " axles which rotate in iron bushings. I have a prime-focus camera attachment, and an electric exhaust fan to control air currents in the tube. Recently I completed a patrol camera, the lens being a 75-mm. $f/4.7$ war-surplus anastigmat. The 8-inch is used as a guide telescope, and I guide by hand since an electric clock drive is still to be completed. With very fast Royal-X Pan cut film (ASA rating 1600), I have secured good star images to the 9th magnitude with a 90-second exposure.

On one side of the hardwood saddle is a 3-inch reflector, which serves as a counterweight for the patrol camera on the other side of the declination axis. When the clock drive is finished, the smaller reflector will serve as a guide telescope for longer exposures with the 8-inch.

Right: Don Parker finds that an old oil drum, formerly used as a support for mirror grinding, makes a convenient observing chair for lunar and planetary observing. Attached to the drum is a basket that holds the eyepiece box, as well as an easel with the clipboard he uses for drawings and making notes.



Below: Before the Parker telescope can be covered with its compact-fitting plywood shelter, the tube must be turned into the meridian and pointed to the north pole of the sky.



The performance of the telescope has been very satisfactory. I have detected, for instance, three of the minor divisions in Saturn's rings, and have observed much detail on Mars, including 11 canals last summer. I have seen 10 spots and craters on the floor of Plato, a lunar feature that I study regularly. My chief interest is in lunar and planetary work.

Moving the telescope's mirror and tube indoors after observing is dangerous and time consuming, so I have constructed the plywood shelter illustrated here. One of the sides is fastened by means of five wing nuts; when these are removed the rest of the weatherproof covering can be easily slid away.

DON PARKER
595 Braeside Rd.
Highland Park, Ill.

SAVE MONEY — USE OUR

Regular Customer Discount Plan

First order, net; second order, 2% discount; third order, 3% discount; fourth order, 4% discount; and on up in savings until

Tenth order — 10% discount

(Orders must be \$25.00 or more. Plan is retro-active to January, 1957.)

We give you:

- PERSONALIZED SERVICE
- PROFESSIONAL QUALITY
- CUSTOM PRODUCTION

New complete 1958 catalogue, 25c (Deductible from first order)

QUALITY OPTICS

Harold E. Snyder, Walbridge, Ohio

OPTICAL FLATS

Pitch Polished, Beral Coated

Rectangular shape

$1\frac{1}{2}$ " x $1\frac{1}{2}$ "
 $\frac{1}{2}$ " wave \$1.75 each
 $\frac{1}{4}$ " wave \$4.50 each
Postpaid.

Elliptical shape

Heavy edges to minimize temperature effects.
 $1\frac{1}{4}$ " minor axis.
 $\frac{1}{2}$ " wave \$4.00 each
 $\frac{1}{4}$ " wave \$6.50 each
Postpaid.

BERAL COATINGS — same optical characteristics as aluminum — mechanically more durable — not over-coated — may be removed without harming glass surface. Prices for Beral coating telescope mirrors: 4" diam. - \$2.75, 6" - \$3.50, 8" - \$4.50, 10" - \$6.50, and $12\frac{1}{2}$ " - \$9.75 each, f.o.b. Skokie.

LEROY M. E. CLAUSING

8038 Monticello Ave.

Skokie, Ill.

TRIGARTH

TURRET

and

Eyepiece

Attachment

with Rack

and Pinion



Just turn the Trigarth Turret and easily improve the performance of your telescope. It holds three eyepieces of standard $1\frac{1}{4}$ " O.D. The Trigarth Turret sells for \$15.95 postpaid. The Eyepiece Attachment with Rack and Pinion also takes standard $1\frac{1}{4}$ " O.D. eyepieces. The rack and pinion is machined from solid aluminum castings, precisely fitted for smooth performance. The main tube is $1\frac{1}{4}$ " long; sliding tube adds 2"; total movement $3\frac{1}{2}$ ". Choice of gray or black crinkle finish. The Eyepiece Attachment with Rack and Pinion is priced at \$15.95 postpaid.

MIRROR CELLS

Made of light, sturdy aluminum, each is ideal for securing the mirror to the tube. The cells are spring adjusted to absorb shocks and are cut away for ventilation.

6" — \$7.00
8" — \$11.50
10" — \$35.00



BUILD YOUR OWN TELESCOPE

Prisms Lenses Eyepieces
Mirrors ground to your order
Aluminizing—with Quartz Coating

Satisfaction Guaranteed

Write for Free Catalog, Instructions, 10¢.

GARTH OPTICAL COMPANY

P. O. Box 991

Springfield 1, Mass.

ORTHO-STAR*

Brand New . . .

Parfocalized . . .

Superb!



OCULARS

ORTHO-STAR* oculars are available in the following focal lengths, giving, for example, the indicated powers when used in conjunction with an 8" f/8 mirror:

27 mm.	61x
20 mm.	81x
16 mm.	102x
10 mm.	163x
7 mm.	246x

Guaranteed to be the finest you ever used — or return for full refund!

Try them at our risk. If you do not agree that the ORTHO-STAR* gives you the clearest and most brilliant image — the finest detail you have ever experienced with your telescope, simply return it within 30 days, in its original condition, for a full cash refund. For more details, see July issue, page 438.

Orders accompanied by payment sent postpaid; if sent C.O.D., you pay postage.

\$19.50 each

*Copyrighted


TRECKER-PATHFINDER


EQUATORIAL MOUNT

\$74.50 complete This mount will accommodate 4-inch to 8-inch telescopes. Specify your tube size when ordering.


Standard 36-inch height — massive 1½-inch steel shafting, in oil-impregnated bronze bearings.

This amazing EQUATORIAL MOUNT is just what the doctor ordered for mounting that homemade telescope you labored so hard to finish. Now you can purchase a beautifully constructed, highly rigid equatorial mount, COMPLETE, for your own telescope as economically as if you had built it yourself. This terrific mount is made entirely of metal; all of the moving equatorial parts are polished to work with maximum ease. Legs, head, and counterweight are all removable for easy storing. The saddle allows complete rotation of your tube. One of the more important features in this mount is that the polar axle is extended for ease in attaching a clock drive and/or setting circles, which may be added at any time. The TRECKER-PATHFINDER mount also has a beautiful, chip-resistant finish. Taking all of these unusual features into consideration, this is truly one of the best DOLLAR-FOR-DOLLAR values ever offered.






RACK-AND-PINION
Focusing Device
\$17.50



SPIDERS
4-vane
6- and 8-inch **\$11.50**
10-inch **\$14.95**
12-inch **\$16.95**

MIRROR CELLS
Skeleton type
6-inch \$6.50
8-inch **\$10.95**
10-inch **\$17.95**
12½-inch **\$21.00**



FOR UNITRON-TYPE REFRACTORS
"FLEX-LINE CONTROL"

Made especially for polar axis of Unitron-type refractors. Provides slow-motion movement from eyepiece position. Specify telescope size when ordering.
3-foot lengths, \$14.95

COAST INSTRUMENT'S PROFESSIONAL MIRROR CLEANING KIT

Protect your mirror — the heart of your reflector — with our Mirror Cleaning Kit. Be astounded at the performance of your present mirror. With newly aluminized mirrors, many years of sparkling viewing can be yours! Ideal for all optics, hard coated or uncoated. **\$2.35 postpaid**

FINDER SCOPE

7x, 50-mm. objective, helical focusing, with mounts and crosshairs. Same as shown on TRECKERSCOPIES.
\$18.50

SMALL EQUATORIAL REFLECTORS COMPLETE

40x — 90x — 120x — 270x 4¼"	\$74.50	60x — 120x 3"	\$29.50
With mount		With mount	

EQUATORIAL MOUNTING PARTS

Due to popular demand, you may now purchase separately any part from either the TRECKER-PATHFINDER or regular TRECKERSCOPE Equatorial Mount, including legs, saddle, equatorial head, column, and so forth. Write for additional information.

Also Available . . .

All types of astronomical books.
Refractor telescopes, new or trade-in.
Mirror making kits, 6- to 12½-inch.
Bushnell binoculars.
Microscopes, \$9.95 to \$400.00.
Low-power hand telescopes.

Write for complete catalogue.

COAST INSTRUMENT, INC.

"IN OPTICS SINCE 1933"



TRECKERSCOPE



MADE IN U. S. A.

RESEARCH QUALITY — MECHANICALLY PRECISE — OPTICALLY PERFECT



10-inch Cassegrain "SKY-GIANT"

STANDARD

6-inch.....\$295.00
8-inch.....\$375.00
10-inch.....\$675.00
12½-inch.....\$995.00

DE LUXE

6-inch.....\$495.00
8-inch.....\$575.00
10-inch.....\$875.00
12½-inch.....\$1150.00

CASSEGRAIN "SKY-GIANT" \$1695.00

All De Luxe models come equipped with clock drive, manual flex-line drive, setting circles, fully rotating tube, and removable casters.



8-inch De Luxe TRECKERSCOPE

Check these features!

- ★ IN-LINE VIEWING.
- ★ IDEAL FOR ASTROPHOTOGRAPHY.
- ★ 1¾" STEEL SHAFTING giving supreme rigidity.
- ★ BRONZE BEARINGS with Teflon inserts.
- ★ OCULARS — your choice of four superb eyepieces plus Goodwin Resolving Power Lens.
- ★ SYNCHRO-SMOOTH RACK-AND-PINION FOCUSING SYSTEM — standard 1¼".
- ★ HIGH POWERS.
- ★ SETTING CIRCLES.
- ★ SLOW-MOTION CONTROLS on both axes.
- ★ FIBERGLASS TUBE.
- ★ SNAP-OUT CASTERS (WHEELS) with vernier adjustment on each leg.
- ★ PHOTOGRAPHIC GUIDE SCOPE — 45x-83x-101x — 2.4" objective, with rack-and-pinion focusing system.
- ★ FINDING SCOPE — 8x 50-mm. objective, helical focusing.
- ★ SUPER-RIGID MOUNTING — massive aluminum castings, 1¼-inch shafts. Surprisingly light — 6-inch or 8-inch scope, about 85 pounds.
- ★ REMOVABLE BALL-BEARING CASTERS — screw-down legs.
- ★ FIBERGLASS FEATHER-LIGHT TUBE.
- ★ FINDING SCOPE — 50-mm. objective — helical focusing.
- ★ SYNCHRO-MESH RACK-AND-PINION FOCUSING SYSTEM.
- ★ SUPERB OPTICS — f/8 parabolic pyrex mirror — guaranteed 1/8 wave and to resolve to Dawes' limit on good nights.
- ★ OCULARS (EYEPIECES) — any three of these eyepieces: 27-mm., 20-mm., 16-mm., 10-mm., 7-mm. — or any two eyepieces and a Goodwin Resolving Power Lens.
- ★ CLOCK DRIVE — standard 110-volt.
- ★ MANUAL FLEX-LINE CONTROL.
- ★ COMPLETE PORTABILITY — assembled or disassembled in minutes.
- ★ EXPOSED FOCAL PLANE — for convenient astrophotography.

• • • USE OUR EXTENDED PAYMENT PLAN ON ALL SCOPES AND MOUNTS • • •

SEND FOR STEREO-VIEWER . . .

Comes with set of 10 color slides showing TRECKERSCOPIES, the TRECKER-PATHFINDER equatorial mount, solar projector, and refractors. **Only 30¢ postpaid**

WRITE FOR OUR NEW FREE PAMPHLET,

"What You Should Know, Look For, And Demand Before Buying Any Telescope."

All prices, unless otherwise indicated, f.o.b. Long Beach, Calif., and subject to change without notice. Nominal crating charge added for all telescopes and mounts. California residents: Add 4% sales tax to all prices.

4811 LONG BEACH BLVD., LONG BEACH 5, CALIF.
Write — Wire — Phone for additional information.

Phone: GARfield 2-3411 or NEVada 6-7683
Dealer Inquiries Invited.

Complete Telescopes



Our Latest
with
Dual Finder
that swings
to either side
of eyepiece.
Find object
with ease,
east or west
of your
meridian.
•
For details and
price list, write

C. C. YOUNG CO.

25 Richard Road, East Hartford 8, Conn.

SKY-GAZERS EXCHANGE

Classified advertising costs 20 cents a word, including address; minimum charge, \$3.00 per ad. Only one for sale ad per issue for each advertiser. Remittance must accompany order. Insertion is guaranteed only on copy received by the 20th of the second month before publication; otherwise, insertion will be made in next issue. We cannot acknowledge classified ads nor the quality of merchandise advertised. Write Ad Dept., *Sky and Telescope*, Harvard Observatory, Cambridge 38, Massachusetts.

FIRST-QUALITY achromatic objective: Main objective from the 13 1/2" M-1 height finder. Clear aperture, 62 mm. (2-7/16"); focal length, 470 mm. (18.5"). Beautifully air-space mounted in 3 1/4" long tube. \$12.50 postpaid. Some matched pairs at \$25.00 per pair. A. Cottone Co., 340 Canal St., New York 13, N. Y.

ALUMINUM TUBING: Sizes 2" through 8", any length. 5¢, \$1.08 per foot. Write for other prices now. Pesca-A, Box 363, Ann Arbor, Mich.

TELESCOPE TUBES: Strong, durable, low cost, any size, any length. Send 3¢ stamp for information. F. Hudnall, 281 Rosslyn Ave., Worthington, Ohio.

TWO TELESCOPE BARGAINS: 12 1/2" f/4, new special order, all-electric Astrola, with 3 eyepieces, cost over \$1,600.00, sacrifice for \$975.00, delivered prepaid. Also 12 1/2" combination Newtonian-Cassegrain, quick-change heads, 7 eyepieces, camera, pendulum-controlled drive; Caltech-Porter design, replacement over \$8,000.00, sacrifice for \$2,500.00, delivered prepaid. Photos available. B. Wack, 2901 Remington Ave., Baltimore 11, Md.

BEGINNER'S Telescope Kit: \$3.00 each, all parts, and 3 lenses. Make 8-power astronomical telescope described in September, 1949. *Sky and Telescope*. Frank Myers, 19200 N. Park Blvd., Shaker Heights, Ohio.

MELLISH 7 1/4" telescope, 100" focal length; objective chipped four places; thumbnail size; 6 eyepieces; tripod pipe mounting; no finder. \$350.00. W. E. Johnson, 42 Sauk Trail, Park Forest, Ill.

METEORITES: Visitors from space. Genuine, guaranteed. Use for research, lectures, and demonstrations. Individually complete, \$1.00, \$3.00, \$6.00, \$10.00, \$15.00. Scientific Laboratory, 2846 Oakley Ave., Baltimore 15, Md.

ASTROLA 8" f/7, clock drive, excellent condition, no oculars, best offer takes it. Bruce Covey, 251 Miller Lane, Rochester 17, N. Y.

REFRACTOR, 3", achromatic, altazimuth. Made by advanced amateur. Brass tube and fittings. Tripod and case. \$98.00 postpaid. Don D'Entremont, 404 Beale St., Wollaston 70, Mass.

RIGID, accurate, low-cost fork scope mounts, 4" to 24", 10" to 24" circles, micro set in right ascension and declination, with drive. Write your needs to Mather Machine Works, Clarksville, Iowa.

PRACTICAL, new "Bibliography for ATM's," second edition, 8 pages, 25¢. Kelvin Masson, 338 Plaza Ave., St. Louis 21, Mo.

SIDEREAL DRIVE for telescope or camera to track stars, accurate simple gearless design, make color transparencies, complete plans, \$1.00. "Space Chart," proportioned planets, sun, moon, and stars, also constellation maps, \$1.00. L. Musgnug, Box 74, Bethel, Conn.

FOR SALE: 2.4" Unitron equatorial, sun screen, Barlow lens, \$190.00. W. L. Wilie, Jr., 1405 McFaddin, Beaumont, Tex.

CELESTIAL CALENDAR

Universal time is used unless otherwise noted.

MINOR PLANET PREDICTIONS

Dembowska, 349, 9.5. November 8, 5:07.5 +29-51; 18, 4:59.8 +30-18; 28, 4:50.2 +30-36. December 8, 4:39.8 +30-42; 18, 4:29.8 +30-37; 28, 4:21.5 +30-26.

Junco, 3, 7.1. November 8, 5:15.3 +0-33; 18, 5:10.8 -0-54; 28, 5:04.0 -1.55. December 8, 4:55.8 -2-24; 18, 4:48.0 -2-16; 28, 4:41.8 -1-33.

Kleopatra, 216, 8.6. November 8, 4:33.0 +12-49; 18, 4:25.8 +10-59; 28, 4:17.5 +9-19. December 8, 4:09.5 +7-57; 18, 4:03.0 +7-01; 28, 3:59.0 +6-30.

ARTIFICIAL-STAR pinholes, .002" in .0015 brass, \$1.00 each. Carl Hand, 1111 Lundvall Ave., Rockford, Ill.

SPECIAL SALE: 8" reflector. Slight defect does not affect optical performance. Needs only base. \$75.00. Duane Allman, 534 South 67th, Milwaukee, Wisc.

COMET SEEKERS: 5" aperture, 20" e.f.l., mounted telescope, \$175.00. Schmidt cameras, any size and focal ratio, typical prices: 7.5" aperture with 10" mirror f/2.8 (optics only), \$400.00; mounted, \$750.00. Arthur De Vany, 1580 S. Ynez Ave., Monterey Park, Calif.

FOR SALE: 8" f/8 mirror, unaluminized, 1/4-wave paraboloid, pyrex. Asking \$55.00. Daniel Kahan, 805 Longshore Ave., Philadelphia 11, Pa.

EXCHANGE VISIT: English amateur, 21, awaiting Cambridge, England, wishes to visit with an American amateur, September, 1958. Return visit later. Write, giving details, to Julian Barbour, South Newington, Nr. Banbury, Oxon, England.

FOR SALE: 2.4" Unitron altazimuth with all standard accessories. Bought last year. \$75.00 f.o.b. Oakland. Carl Backlund, 1200 Alpine Rd., Apt. 50, Walnut Creek, Calif.

BARDON TELESCOPE: 2 1/2" clear aperture, 3 eyepieces, altazimuth mounting; owned by astronomy club. Send bids over \$110.00 or write for information to Rasmussen and Reece, Amsterdam, N. Y.

FOR SALE: 7 x 50 binocular eyepieces, spiral focusing, coated eye lens, \$3.00 each. Height finder objectives, 2-3/64" diameter, coated, achromatic, 9" focal length, in threaded metal cell, \$5.00 each. Prisms, right-angle, 3/4" x 3/4" face, 1-5/16" hypotenuse, \$1.00 each. Tank periscope prisms, right angle, 2 1/4" hypotenuse, 5 1/4" long in metal and plastic housing, two for \$1.50. All postpaid. Satisfaction or money refunded. Charles E. Roberts, 730 4th St., Fremont, Ohio.

PHOTOGRAPH NEEDED: Stars over sea, no land. Will buy for publication. Science Editor, Artists and Writers Guild, 630 Fifth Ave., New York 20, N. Y.

REFLECTOR, 8", beral coated, cast-iron equatorial mount, synchronous drive, finder, 3 eyepieces, catter dolly. \$185.00, plus transportation. C. Johnson, 48 Roberts St., Watertown, Conn.

NOW! You can see with both eyes while observing! Transform your telescope to get the results of giant binoculars by using the Brayton Binocular Eyepiece Holder (see page 607, October issue of *Sky and Telescope*). Brochure on request. Brayton Optical Exchange, 120 Beachway, Monterey, Calif.

INTERESTED in astronomy as a career? *Vocational and Professional Monographs: Astronomy* by Freeman D. Miller describes personal qualifications, scholastic training, and job opportunities. \$1.00 postpaid. Send to Box B, *Sky and Telescope*, Harvard Observatory, Cambridge 38, Mass.

WANTED: Volumes I through IX of *Sky and Telescope*. State numbers available, price, and condition. February and November, 1952, issues wanted also. James Brown, 1409 Belmead Drive, Kingsport, Tenn.

CANADIAN AMATEURS: America's best-selling line of refracting telescopes and accessories, now available from Canada's largest optical specialists. Save customs clearance and extra duty. Carsten Instruments, Ltd., 88 Lycos Drive, Toronto 10, Canada.

SCHMIDT'S famous moon map, 25 plates. Printed on 8 1/2" x 11" sheets. \$1.75 postpaid world-wide. Scientific Book Service, 663 Franklin Ave., Columbus 15, Ohio.

SOLAR FURNACE, 36", mounted, 14" focal length. Packed and shipped, \$125.00. Stanley R. Armitage, 4813 Jonestown Rd., Harrisburg, Pa.

Melpomene, 18, 8.3. November 8, 4:57.7 +2-42; 18, 4:50.1 +1-55; 28, 4:40.5 +1-34. December 8, 4:30.5 +1-45; 18, 4:21.9 +2-26; 28, 4:15.9 +3-32.

After the asteroid's name are its number and the magnitude expected at opposition. At 10-day intervals are given its right ascension and declination (1950.0) for 0^h Universal time. In each case the motion of the asteroid is retrograde. Data are supplied by the IAU Minor Planet Center at the University of Cincinnati Observatory.

VARIABLE STAR MAXIMA

November 2, RS Herculis, 171723, 8.0; 2, T Normae, 153654, 7.4; 3, Omicron Ceti (Mira), 021403, 3.7; 4, R Normae, 152849, 7.2; 6, R Indi, 222867, 8.0; 18, S Gruis, 221948, 7.8; 18, T Centauri, 133633, 6.1; 29, R Leonis Minoris, 093934, 7.2.

December 2, R Andromedae, 001838, 7.0; 2, T Ursae Majoris, 123160, 7.9; 5, R Draconis, 163266, 7.6; 5, R Horologii, 025050, 6.0; 8, S Canis Minoris, 072708, 7.5; 9, RR Scorpii, 165030, 6.0.

These predictions of variable star maxima are by the AA.VSO. Only stars are included whose mean maximum magnitudes are brighter than magnitude 8.0. Some, but not all of them, are nearly as bright as maximum two or three weeks before and after the dates for maximum. The data given include, in order, the day of the month near which the maximum should occur, the star name, the star designation number, which gives the rough right ascension (first four figures) and declination (bold face if southern), and the predicted magnitude.

OCCULTATION PREDICTIONS

November 8-9 Epsilon Tauri 3.6, 4:26.1 +19-05.2, 17. Im: F 10:27.0 -1.9 +0.1 69; H 9:42.0 -2.2 +1.5 56. Em: F 11:37.9 -0.8 -1.9 294; H 10:58.8 -1.9 -2.0 296.

November 11-12 Lambda Geminorum 3.6, 7:15.6 +16-37.2, 20. Im: B 11:10.3 -0.6 -2.8 146; D 11:13.0 -0.4 -3.9 162.

For stations in the United States and Canada, usually for stars of magnitude 5.0 or brighter, data from the *American Ephemeris* and the *British Nautical Almanac* are given here, as follows: evening-morning date, star name, magnitude, right ascension in hours and minutes, declination in degrees and minutes, moon's age in days, immersion or emersion; standard station designation, UT, a and b quantities in minutes, position angle on the moon's limb; the same data for each standard station westward.


The a and b quantities tabulated in each case are variations of standard-station predicted times per degree of longitude and of latitude, respectively, enabling computation of fairly accurate times for one's local station (long. Lo, lat. L) within 200 or 300 miles of a standard station (long. LoS, lat. LS). Multiply a by the difference in longitude (Lo - LoS), and multiply b by the difference in latitude (L - LS), with due regard to arithmetic signs, and add both results to (or subtract from, as the case may be) the standard-station predicted time to obtain time at the local station. Then convert the Universal time to your standard time.

Longitudes and latitudes of standard stations are:
A +72° 5', +42° 5' E +91° 0', +40° 0'
B +73° 6', +45° 5' F +98° 0', +31° 0'
C +77° 1', +38° 9' G Discontinued
D +79° 4', +43° 7' H +120° 0', +36° 0'
I +123° 1', +49° 5'

MINIMA OF ALGOL

November 2, 10:05; 5, 6:53; 8, 3:42; 11, 0:31; 13, 21:20; 16, 18:09; 19, 14:58; 22, 11:47; 25, 8:36; 28, 5:25. December 1, 2:14; 3, 23:03; 6, 19:52; 9, 16:41.

These minima predictions for Algol are based on the formula in the 1953 *International Supplement of the Krakow Observatory*. The times given are geocentric; they can be compared directly with observed times of least brightness.



THE SUN, MOON, AND PLANETS THIS MONTH

The sun, on the ecliptic, is shown for the beginning and end of the month. The moon's symbols give its phase roughly, with the date marked alongside. Each planet is located for the middle of the month or for other dates shown. All positions are for 0^h Universal time on the respective dates.

Mercury will be in the evening sky all month, but poorly situated for viewing. It may be observed during the last few days of November as a -0.3-magnitude object, setting one hour after the sun.

Venus' angular distance from the sun attains a maximum of 47° 14' on November 18th, the planet then being at greatest eastern elongation. On that date it will be of magnitude -4.0, a brilliant white object in Sagittarius setting in the southwestern sky three hours after the sun. Telescopically, Venus will present a half-illuminated disk 25.1 seconds of arc in diameter.

There will be a close conjunction with the 6.5-magnitude star 86B Sagittarii on the 11th, but the minimum separation of about one diameter of the planet will be at 13:11 UT, during daylight hours for the Americas.

The moon will be totally eclipsed on November 7th as seen from eastern Asia, the Pacific, and the western part of North America. For details see page 21 of this issue.

Mars, now 2nd magnitude, rises two hours before the sun by the end of the month. The reddish planet is traveling

rapidly eastward from Virgo into Libra. A very close conjunction with Lambda Virginis, magnitude 4.6, occurs on November 18th at 8:59 UT, Mars passing only about four seconds of arc from the star. Closest approach will take place before Mars rises for observers in the United States.

Jupiter is viewed rising in the east nearly three hours ahead of the sun in midmonth. The giant yellow planet is moving eastward in Virgo, and will be 3½° north of Spica on the 21st.

Saturn may be seen only early in November, low in the southwestern sky. On the 7th it sets 1½ hours after sundown, appearing of magnitude +0.7.

Uranus rises about midnight, passing western quadrature with the sun on November 4th. The planet's position among the stars changes very little, because retrograde motion begins on the 17th. Uranus can be found in binoculars as a 6th-magnitude object about 4° east of Delta Cancri.

Neptune, in the morning sky, is too near the sun to be seen this month.

MOON PHASES AND DISTANCE

Full moon	November 7, 14:32
Last quarter	November 14, 21:59
New moon	November 21, 16:19
First quarter	November 29, 6:57
Full moon	December 7, 6:16

	November	Distance	Diameter
Apogee	2, 12h	251,700 mi.	29' 30"
Perigee	18, 11h	227,700 mi.	32' 37"
Apogee	30, 7h	251,300 mi.	29' 33"

	December	Distance	Diameter
Perigee	14, 5h	230,100 mi.	32' 16"

UNIVERSAL TIME (UT)

TIMES used in Celestial Calendar are Greenwich civil or Universal time, unless otherwise noted. This is 24-hour time, from midnight to midnight; times greater than 12:00 are p.m. Subtract the following hours to convert to standard times in the United States: EST, 5; CST, 6; MST, 7; PST, 8. If necessary, add 24 hours to the UT before subtracting, in which case the result is your standard time on the day preceding the Greenwich date shown.

12" Sky Master



The most practical way really to learn **WHAT** can be seen in the sky, and **WHEN**, and **WHERE** at a glance!

The Sky Master "functional globe" shows the sky exactly as you see it, and not backwards. Each of the 12 segments is dated for a 9 p.m. view. Shows all stars to magnitude 5.5, nebulae, galactic and ecliptic poles, and the like. The 57 navigation stars have the official star symbols, numbers; geometrical constellation figures.

Invaluable for students of General Astronomy, and of Marine and Air Navigation, as co-ordinates are given in RAH, SHA, and LHA-Aries.

Guide and Check Lists

Introduce Astronomy in one night!

Guide: a complete, up-to-date, short course in astronomy and navigation; with diagrams.

Main List: 550 entries in 20 columns, with both 1960 SHA and RAH co-ordinates. Stars to magnitude 3.5; variable and multiple stars; star colors, distances, and both magnitudes; nebulae, radiants, and other data.

15 Lists for quickly locating sky objects. Complete solar system data. Monthly planetary positions to 1965. Constellations and 190 star names and their meanings, 108 brightest and 50 largest stars. Messier objects and NGC-number cross lists. LHA-Aries on meridian table, with sunrise, sunset, and twilight columns; and others.

12" globe "do-it-yourself" kit \$5.00

Guide and Check Lists, with above order 1.75

Guide and Check Lists, separately 1.95

12" globe, assembled (with above book and plain wire stand) f.o.b. 14.00

24" globe for schools; prices sent upon request.

Globes may be had with, or without, the mythological pictures (printed in blue).

SKY MASTER, Capt. J. Ellrich

781 Fairview Ave. Fairview, New Jersey

Dyno-Speed

WIND INDICATOR

First Time in Kit Form

Now available in kit form at amazing low price. Easily assembled in 30 to 40 min. with screwdriver and pliers. Each instrument is skillfully engineered for fine performance and long life, and comes complete with 0 to 100 M.P.H. indicator. Kit also includes mounting ideal for boats, homes, airports, weather stations, etc. Instrument is self generating—needs no batteries.

Send Check or Money Order Sorry, no C.O.D.'s


Write for free literature.

DYNO-SPEED INSTRUMENT CO.

P.O. Box 47 Lukewood, N. J.



COMPLETE \$21.95 (P.O.B. Lukewood)




SOUTHERN STARS

The sky as seen from latitudes 20° to 40° south, at 11 p.m. and 10 p.m., local time, on the 7th and 23rd of January, re-

spectively; also at 9 p.m. and 8 p.m. on February 6th and 21st. For other dates, add or subtract $\frac{1}{2}$ hour per week.

Speaking of star-people appearing upside down in the Southern Hemisphere,

on this chart some very famous ones are now seen that way: Orion, Auriga, Perseus, while even the Twins of Gemini have their feet higher than their heads, the stars Castor and Pollux.



STARS FOR NOVEMBER

The sky as seen from latitudes 30° to 50° north, at 9 p.m. and 8 p.m., local time, on the 7th and 23rd of November,

respectively; also, at 7 p.m. and 6 p.m. on December 7th and 23rd. For other dates, add or subtract ½ hour per week.

As the large bird constellations of our northern summer, Aquila the Eagle and

Cygnus the Swan, fly down the western evening sky, two giant animals, Taurus the Bull and Cetus, monster of the deep, dominate the heavens in the east, pursued by Orion the Hunter.

NOW — THE DELUXE 4" DYNASCOPE Reflector

with ADVANCED Precision Features Offers

You a Truly Amazing Telescope COMPLETE

For Only **\$79⁹⁵**

f.o.b. Hartford, Conn.
Shipping Wt. 21 lbs.
Express charges collect



Inquire for details of convenient Time-Payment Plan.

Compare these advanced features with any telescope at double the price!

- (1) **FOUR-INCH PARABOLOIDAL PYREX MIRROR** — Finished to exacting specifications and guaranteed to perform to Dawes' limit for this size of instrument. Aluminized and zircon-quartz overlaid to insure maximum protection and lasting use. The 4-inch mirror gathers one third more light than a 3½-inch mirror.
- (2) **IMPROVED EQUATORIAL MOUNTING** — Rugged cast iron, weighing about 12 pounds, guaranteed vibration free. Wing clamp shown in the inset provides easy change of latitude setting if you move to a new observing station. Free-moving polar and declination axles are ⅝-inch steel, supported on four bearing surfaces, guaranteed to have necessary smoothness in operation. Friction clamp for declination and large knurled brass adjustment for right ascension.
- (3) **EYEPIECE MOUNT** — Standard 1¼ inch with exclusive double-draw and rack-and-pinion focusing. Adjustable for three inches of travel to accommodate any eyepiece, positive or negative. Built-in diagonal mirror has a surface finished to ⅛ wave.
- (4) **THREE EYEPIECES** — 18-mm. Huygens, 9-mm. achromatic Ramsden, 7-mm. achromatic Ramsden, giving powers of 65x, 130x, and 167x.
- (5) **ACHROMATIC FINDER** — 4-power with crosshairs; extra large field of view.
- (6) **COVERS** for eyepiece tube and open end of the telescope itself.
- (7) **BAKELITE TUBE** beautifully finished in gray wrinkle enamel.
- (8) **IMPROVED TRIPOD** — Natural-finish hardwood legs, folding against each other. Chain ties provided (not illustrated). Sturdy, balanced, providing perfect portability.

THE SUPERIOR 4-INCH TELESCOPE. *Prove it yourself at NO RISK!*

Here is the scientific instrument that serious amateurs have been waiting for — with a full warranty of high performance at low cost!

Now you need not spend \$150 and up to be sure of excellent observing. Nor do you need to go to the time and trouble of building your own telescope to get the most value for your money. For the dollars-and-cents facts about the Deluxe Dynascope are these:

The advanced precision features are those you would select for yourself. Buying them singly, as an individual, you could never beat our low price. You could not hope to surpass the mechanical excellence and ruggedness that have been engineered into this superb instrument to meet the most exacting standards! Our instrument has been tested and approved by staff members of leading planetariums.

The Deluxe Dynascope comes to you complete with every part and feature exactly as described and illustrated here. Each instrument is carefully triple-tested before shipment and is accompanied by the inspector's

performance report. Specially packed, it is ready to be set up for observing a few minutes after unpacking. Shipment is f.o.b. Hartford, Conn., express charges collect (weight 21 pounds). There is nothing else to buy, no added charges, no extras of any kind needed.

YOU WILL BE DELIGHTED—OR MONEY BACK!

Prove to yourself — without risk — how good the Deluxe Dynascope really is. Order it now. Try it at your own pleasure. Compare its performance with that of any other telescope at double the cost or more. It will delight you and exceed your every expectation — or simply return it within 30 days for a full refund. But don't delay. As you know, quality engineering of precision instruments does not permit mass production. Consequently the supply is limited. To assure yourself of an early delivery, send your check or money order today!


CRITERION MANUFACTURING COMPANY

Manufacturers of Quality Optical Instruments

DEPT. STD-41, 331 CHURCH ST., HARTFORD 1, CONN. TEL.: CHAPEL 7-1696 — CABLE ADDRESS: CRICO

EXPERTS IN DOMES AND TELESCOPES

ASTRO-DOME, expert in the construction of observatory domes, is pleased to announce the completion of an Astro-Observatory for the Reed telescope at Yale University, New Haven, Connecticut. This building is 16 feet 6 inches in diameter and weighs 13,000 pounds. The dome is fabricated from heavy structural steel members and is covered with 16-gauge galvanized-steel sheets. The result is a dome offering smooth appearance plus great strength. The Astro-Observatory illustrated here may be obtained in diameters up to 50 feet. May we assist you with your installation?



TINSLEY LABORATORIES, expert in the production of precision instruments, thoroughly tests every optical system for absolute conformance to the highest specifications. All optical surfaces are polished to 1/10-wave accuracy, and each system is guaranteed to reach theoretical limits of resolution. Through research, development, and construction of optical devices for the United States government and industrial organizations, Tinsley Laboratories is able to furnish your telescope at lower cost without sacrifice in quality. Small and large telescopes of any design are available to the required precision. You are invited to request information of any kind that would be useful to you.

As experts in domes and telescopes, Astro-Dome and Tinsley Laboratories now make possible a complete observatory from telescope to housing at a cost that will be pleasantly reasonable. Write either company for details, which will be furnished without obligation.

ASTRO-DOME MANUFACTURING, INC.


Box 217, Station A, Canton 5, Ohio
Telephone: GLEndale 4-2755



Tinsley
laboratories

2530 Grove St., Berkeley 4, Calif.

UNITRON



It takes only \$12.50 down to own this UNITRON

Little wonder that the UNITRON 2.4" Altazimuth Refractor has long been America's most popular low-priced refractor.

For UNITRON and only UNITRON combines unexcelled optical and mechanical performance with exclusive features designed to increase your observing pleasure. As an example, take UNIHEX — the Rotary Eyepiece Selector which relieves you of the burden of fumbling with eyepieces in the dark. Or, perhaps your observing situation may call for the DUETRON Double Eyepiece, the erecting prism system, the Astro-Camera 220, or the sun screen.

Optically speaking, UNITRON Refractors duplicate the performance of larger telescopes of other types. Furthermore, there are no mirror surfaces to become tarnished,

no secondary optics to cause diffraction patterns, and no folding of the light back on itself with consequent loss of definition. It is not surprising that you see more and see better with a UNITRON.

The UNITRON 2.4" Altazimuth Refractor is priced at only \$125 complete with tripod, sturdy mounting with micrometric slow-motion controls, four eyepieces, carrying cases, and accessories. Using our Easy Payment Plan, a mere 10% down payment of only \$12.50 brings this UNITRON to you.

Interested? It is really not too early to let Santa know how you feel.

See pages 28 and 29.

UNITRON

INSTRUMENT DIVISION of UNITED SCIENTIFIC CO.
204-206 MILK STREET BOSTON 9, MASSACHUSETTS

